


JEE April 2019

Test Date	12/04/2019
Test Time	2:30 PM - 5:30 PM
Subject	Paper I EH

Section : Physics

- Q.1** A transparent cube of side d , made of a material of refractive index μ_2 , is immersed in a liquid of refractive index μ_1 ($\mu_1 < \mu_2$). A ray is incident on the face AB at an angle θ (shown in the figure). Total internal reflection takes place at point E on the face BC.



Then θ must satisfy :

Options

1. $\theta < \sin^{-1} \sqrt{\frac{\mu_2^2}{\mu_1^2} - 1}$ ✓
2. $\theta > \sin^{-1} \frac{\mu_1}{\mu_2}$
3. $\theta < \sin^{-1} \frac{\mu_1}{\mu_2}$
4. $\theta > \sin^{-1} \sqrt{\frac{\mu_2^2}{\mu_1^2} - 1}$

Question Type : **MCQ**

Question ID : **41652913437**

Option 1 ID : **41652952526**

Option 2 ID : **41652952528**

Option 3 ID : **41652952529**

Option 4 ID : **41652952527**

Status : **Answered**

Chosen Option : **2**

Q.2 A Carnot engine has an efficiency of $1/6$.
When the temperature of the sink is reduced by 62°C , its efficiency is doubled.
The temperatures of the source and the sink are, respectively,

- Options
1. 124°C , 62°C
 2. 37°C , 99°C ✓
 3. 99°C , 37°C
 4. 62°C , 124°C

Question Type : **MCQ**

Question ID : **41652913426**

Option 1 ID : **41652952483**

Option 2 ID : **41652952484**

Option 3 ID : **41652952482**

Option 4 ID : **41652952485**

Status : **Not Answered**

Chosen Option : --

Q.3 A diatomic gas with rigid molecules does 10 J of work when expanded at constant pressure. What would be the heat energy absorbed by the gas, in this process ?

- Options
1. 40 J
 2. 35 J ✓
 3. 25 J
 4. 30 J

Question Type : **MCQ**

Question ID : **41652913427**

Option 1 ID : **41652952489**

Option 2 ID : **41652952488**

Option 3 ID : **41652952486**


Option 4 ID : **41652952487**

Status : **Answered**

Chosen Option : **2**

Q.4

A smooth wire of length $2\pi r$ is bent into a circle and kept in a vertical plane. A bead can slide smoothly on the wire. When the circle is rotating with angular speed ω about the vertical diameter AB, as shown in figure, the bead is at rest with respect to the circular ring at position P as shown. Then the value of ω^2 is equal to :



- Options
1. $2g/r$
 2. $\frac{\sqrt{3}g}{2r}$
 3. $2g/(r\sqrt{3})$ ✓
 4. $(g\sqrt{3})/r$

Question Type : **MCQ**

Question ID : **41652913421**

Option 1 ID : **41652952462**

Option 2 ID : **41652952463**

Option 3 ID : **41652952465**

Option 4 ID : **41652952464**

Status : **Not Answered**

Chosen Option : --

Q.5 Let a total charge $2Q$ be distributed in a sphere of radius R , with the charge density given by $\rho(r) = kr$, where r is the distance from the centre. Two charges A and B, of $-Q$ each, are placed on diametrically opposite points, at equal distance, a , from the centre. If A and B do not experience any force, then :

- Options
1. $a = \frac{3R}{2^{1/4}}$

2. $a = R/\sqrt{3}$
3. $a = 2^{-1/4} R$
4. $a = 8^{-1/4} R$ ✓

Question Type : **MCQ**

Question ID : **41652913430**

Option 1 ID : **41652952501**

Option 2 ID : **41652952500**

Option 3 ID : **41652952499**

Option 4 ID : **41652952498**

Status : **Answered**

Chosen Option : **2**

Q.6 The electron in a hydrogen atom first jumps from the third excited state to the second excited state and subsequently to the first excited state. The ratio of the respective wavelengths, λ_1/λ_2 , of the photons emitted in this process is :

Options 1. $7/5$

2. $20/7$ ✓

3. $9/7$

4. $27/5$

Question Type : **MCQ**

Question ID : **41652913440**

Option 1 ID : **41652952539**

Option 2 ID : **41652952541**

Option 3 ID : **41652952540**


Option 4 ID : **41652952538**

Status : **Answered**

Chosen Option : **2**

Q.7

A block of mass 5 kg is (i) pushed in case (A) and (ii) pulled in case (B), by a force $F = 20 \text{ N}$, making an angle of 30° with the horizontal, as shown in the figures. The coefficient of friction between the block and floor is $\mu = 0.2$. The difference between the accelerations of the block, in case (B) and case (A) will be : ($g = 10 \text{ ms}^{-2}$)



- Options
1. 3.2 ms^{-2}
 2. 0 ms^{-2}
 3. 0.8 ms^{-2} ✓
 4. 0.4 ms^{-2}

Question Type : **MCQ**

Question ID : **41652913419**

Option 1 ID : **41652952455**

Option 2 ID : **41652952454**

Option 3 ID : **41652952457**

Option 4 ID : **41652952456**

Status : **Answered**

Chosen Option : **3**

Q.8 A small speaker delivers 2 W of audio output. At what distance from the speaker will one detect 120 dB intensity sound ?
[Given reference intensity of sound as 10^{-12} W/m^2]

- Options
1. 40 cm ✓
 2. 20 cm
 3. 10 cm
 4. 30 cm

Question Type : **MCQ**

Question ID : **41652913428**

Option 1 ID : **41652952493**

Option 2 ID : **41652952491**

Option 3 ID : **41652952490**

Option 4 ID : **41652952492**

Status : **Not Answered**

Chosen Option : --

Q.9 Consider an electron in a hydrogen atom, revolving in its second excited state (having radius 4.65 \AA). The de-Broglie wavelength of this electron is :

- Options
1. 12.9 \AA
 2. 6.6 \AA
 3. 9.7 \AA ✓
 4. 3.5 \AA

Question Type : MCQ

Question ID : 41652913439

Option 1 ID : 41652952537

Option 2 ID : 41652952535


Option 3 ID : 41652952536

Option 4 ID : 41652952534

Status : Not Answered

Chosen Option : --

Q.10 Figure shows a DC voltage regulator circuit, with a Zener diode of breakdown voltage = 6V . If the unregulated input voltage varies between 10V to 16V , then what is the maximum Zener current ?



- Options
1. 1.5 mA
 2. 3.5 mA ✓
 3. 7.5 mA
 4. 2.5 mA

Question Type : MCQ

Question ID : 41652913442

Option 1 ID : 41652952547

Option 2 ID : 41652952549


Option 3 ID : 41652952546

Option 4 ID : 41652952548

Status : Not Answered

Chosen Option : --

- Q.11 An electron, moving along the x -axis with an initial energy of 100 eV, enters a region of magnetic field $\vec{B} = (1.5 \times 10^{-3} \text{T}) \hat{k}$ at S (See figure). The field extends between $x=0$ and $x=2$ cm. The electron is detected at the point Q on a screen placed 8 cm away from the point S. The distance d between P and Q (on the screen) is :
(electron's charge = $1.6 \times 10^{-19} \text{C}$, mass of electron = $9.1 \times 10^{-31} \text{kg}$)



- Options
1. 1.22 cm
 2. 12.87 cm ✓
 3. 11.65 cm
 4. 2.25 cm

Question Type : MCQ

Question ID : 41652913433

Option 1 ID : 41652952510

Option 2 ID : 41652952511

Option 3 ID : 41652952512

Option 4 ID : 41652952513

Status : Not Answered

Chosen Option : --

Q.12

A tuning fork of frequency 480 Hz is used in an experiment for measuring speed of sound (v) in air by resonance tube method. Resonance is observed to occur at two successive lengths of the air column, $l_1 = 30$ cm and $l_2 = 70$ cm. Then, v is equal to :

- Options
1. 384 ms^{-1} ✓
 2. 332 ms^{-1}
 3. 338 ms^{-1}
 4. 379 ms^{-1}

Question Type : **MCQ**

Question ID : **41652913444**

Option 1 ID : **41652952556**

Option 2 ID : **41652952557**

Option 3 ID : **41652952554**

Option 4 ID : **41652952555**

Status : **Answered**

Chosen Option : 1

Q.13 A spring whose unstretched length is l has a force constant k . The spring is cut into two pieces of unstretched lengths l_1 and l_2 where, $l_1 = nl_2$ and n is an integer. The ratio k_1/k_2 of the corresponding force constants, k_1 and k_2 will be :

- Options
1. $\frac{1}{n^2}$
 2. n^2
 3. n
 4. $\frac{1}{n}$ ✓

Question Type : **MCQ**

Question ID : **41652913420**

Option 1 ID : **41652952461**

Option 2 ID : **41652952460**

Option 3 ID : **41652952458**

Option 4 ID : **41652952459**

Status : **Not Answered**

Chosen Option : --

Q.14

Two sources of sound S_1 and S_2 produce sound waves of same frequency 660 Hz. A listener is moving from source S_1 towards S_2 with a constant speed u m/s and he hears 10 beats/s. The velocity of sound is 330 m/s. Then, u equals :

- Options
1. 10.0 m/s
 2. 2.5 m/s ✓
 3. 5.5 m/s
 4. 15.0 m/s

Question Type : **MCQ**

Question ID : **41652913429**

Option 1 ID : **41652952496**

Option 2 ID : **41652952494**

Option 3 ID : **41652952495**

Option 4 ID : **41652952497**

Status : **Not Answered**

Chosen Option : --

Q.15 A moving coil galvanometer, having a resistance G , produces full scale deflection when a current I_g flows through it. This galvanometer can be converted into (i) an ammeter of range 0 to I_0 ($I_0 > I_g$) by connecting a shunt resistance R_A to it and (ii) into a voltmeter of range 0 to V ($V = GI_0$) by connecting a series resistance R_V to it. Then,

Options

$$R_A R_V = G^2 \left(\frac{I_0 - I_g}{I_g} \right) \text{ and}$$

1.

$$\frac{R_A}{R_V} = \left(\frac{I_g}{I_0 - I_g} \right)^2$$

2. $R_A R_V = G^2$ and $\frac{R_A}{R_V} = \frac{I_g}{I_0 - I_g}$

$$R_A R_V = G^2 \left(\frac{I_g}{I_0 - I_g} \right) \text{ and}$$

3.

$$\frac{R_A}{R_V} = \left(\frac{I_0 - I_g}{I_g} \right)^2$$

4. $R_A R_V = G^2$ and $\frac{R_A}{R_V} = \left(\frac{I_g}{I_0 - I_g} \right)^2$ ✓

Question Type : **MCQ**

Question ID : **41652913445**

Option 1 ID : **41652952561**

Option 2 ID : **41652952559**

Option 3 ID : **41652952560**

Option 4 ID : **41652952558**

Status : **Not Answered**

Chosen Option : --

Q.16 In an amplitude modulator circuit, the carrier wave is given by,
 $C(t) = 4 \sin(20000 \pi t)$ while modulating signal is given by, $m(t) = 2 \sin(2000 \pi t)$. The values of modulation index and lower side band frequency are :

- Options
1. 0.5 and 9 kHz ✓
 2. 0.4 and 10 kHz
 3. 0.5 and 10 kHz
 4. 0.3 and 9 kHz

Question Type : **MCQ**

Question ID : **41652913443**

Option 1 ID : **41652952553**

Option 2 ID : **41652952551**

Option 3 ID : **41652952552**

Option 4 ID : **41652952550**

Status : **Not Answered**

Chosen Option : --

Q.17 A particle is moving with speed $v = b\sqrt{x}$ along positive x -axis. Calculate the speed of the particle at time $t = \tau$ (assume that the particle is at origin at $t = 0$).

- Options
1. $b^2 \tau$
 2. $\frac{b^2 \tau}{\sqrt{2}}$
 3. $\frac{b^2 \tau}{2}$ ✓
 4. $\frac{b^2 \tau}{4}$

Question Type : **MCQ**Question ID : **41652913417**Option 1 ID : **41652952449**Option 2 ID : **41652952447**Option 3 ID : **41652952448**Option 4 ID : **41652952446**Status : **Answered**Chosen Option : **4**

Q.18 The ratio of the weights of a body on the Earth's surface to that on the surface of a planet is 9 : 4. The mass of the planet is $\frac{1}{9}$ th of that of the Earth. If 'R' is the radius of the Earth, what is the radius of the planet ? (Take the planets to have the same mass density)

Options

1. $\frac{R}{4}$

2. $\frac{R}{2}$ ✓

3. $\frac{R}{3}$

4. $\frac{R}{9}$

Question Type : **MCQ**Question ID : **41652913423**Option 1 ID : **41652952473**Option 2 ID : **41652952470**Option 3 ID : **41652952471**Option 4 ID : **41652952472**Status : **Answered**Chosen Option : **2**

Q.19 The number density of molecules of a gas depends on their distance r from the origin as, $n(r) = n_0 e^{-\alpha r^4}$. Then the total number of molecules is proportional to :

Options 1. $n_0 \alpha^{-3}$

2. $\sqrt{n_0} \alpha^{1/2}$

3. $n_0 \alpha^{-3/4}$ ✓

4. $n_0 \alpha^{1/4}$

Question Type : **MCQ**

Question ID : 41652913416

Option 1 ID : 41652952442

Option 2 ID : 41652952443

Option 3 ID : 41652952444

Option 4 ID : 41652952445

Status : Not Answered

Chosen Option : --

Q.20 A uniform cylindrical rod of length L and radius r , is made from a material whose Young's modulus of Elasticity equals Y . When this rod is heated by temperature T and simultaneously subjected to a net longitudinal compressional force F , its length remains unchanged. The coefficient of volume expansion, of the material of the rod, is (nearly) equal to :

- Options
1. $F/(3\pi r^2 Y T)$
 2. $9F/(\pi r^2 Y T)$
 3. $6F/(\pi r^2 Y T)$
 4. $3F/(\pi r^2 Y T)$ ✓

Question Type : MCQ

Question ID : 41652913424

Option 1 ID : 41652952477

Option 2 ID : 41652952475

Option 3 ID : 41652952474

Option 4 ID : 41652952476


Status : Answered

Chosen Option : 2

Q.21

Find the magnetic field at point P due to a straight line segment AB of length 6 cm carrying a current of 5 A. (See figure)

$$(\mu_0 = 4\pi \times 10^{-7} \text{ N-A}^{-2})$$



- Options
1. $1.5 \times 10^{-5} \text{ T}$ ✓
 2. $2.0 \times 10^{-5} \text{ T}$
 3. $3.0 \times 10^{-5} \text{ T}$
 4. $2.5 \times 10^{-5} \text{ T}$

Question Type : MCQ

Question ID : 41652913434

Option 1 ID : 41652952514

Option 2 ID : 41652952515


Option 3 ID : 41652952517

Option 4 ID : 41652952516

Status : Answered

Chosen Option : 1

- Q.22 Consider the LR circuit shown in the figure. If the switch S is closed at $t=0$ then the amount of charge that passes through the battery between $t=0$ and $t=\frac{L}{R}$ is :



- Options
1. $\frac{7.3 EL}{R^2}$
 2. $\frac{EL}{7.3R^2}$

3. $\frac{2.7 EL}{R^2}$

4. $\frac{EL}{2.7R^2}$ ✓

Question Type : **MCQ**

Question ID : **41652913435**

Option 1 ID : **41652952518**

Option 2 ID : **41652952521**


Option 3 ID : **41652952519**

Option 4 ID : **41652952520**

Status : **Not Answered**

Chosen Option : --

- Q.23** Three particles of masses 50 g, 100 g and 150 g are placed at the vertices of an equilateral triangle of side 1 m (as shown in the figure). The (x, y) coordinates of the centre of mass will be :



Options

1. $\left(\frac{7}{12} \text{ m}, \frac{\sqrt{3}}{4} \text{ m}\right)$ ✓

2. $\left(\frac{7}{12} \text{ m}, \frac{\sqrt{3}}{8} \text{ m}\right)$

3. $\left(\frac{\sqrt{3}}{4} \text{ m}, \frac{5}{12} \text{ m}\right)$

4. $\left(\frac{\sqrt{3}}{8} \text{ m}, \frac{7}{12} \text{ m}\right)$

Question Type : **MCQ**

Question ID : **41652913422**

Option 1 ID : **41652952467**

Option 2 ID : **41652952466**

Option 3 ID : **41652952468**

Option 4 ID : **41652952469**

Status : **Answered**

Chosen Option : 1

Q.24 Half lives of two radioactive nuclei A and B are 10 minutes and 20 minutes, respectively. If, initially a sample has equal number of nuclei, then after 60 minutes, the ratio of decayed numbers of nuclei A and B will be :

- Options
1. 3 : 8
 2. 8 : 1
 3. 1 : 8
 4. 9 : 8 ✓

Question Type : **MCQ**

Question ID : **41652913441**

Option 1 ID : **41652952545**

Option 2 ID : **41652952544**

Option 3 ID : **41652952542**

Option 4 ID : **41652952543**

Status : **Answered**

Chosen Option : **3**

Q.25 Two particles are projected from the same point with the same speed u such that they have the same range R , but different maximum heights, h_1 and h_2 . Which of the following is correct ?

- Options
1. $R^2 = h_1 h_2$
 2. $R^2 = 4 h_1 h_2$
 3. $R^2 = 2 h_1 h_2$
 4. $R^2 = 16 h_1 h_2$ ✓

Question Type : **MCQ**

Question ID : **41652913418**

Option 1 ID : **41652952453**

Option 2 ID : **41652952451**

Option 3 ID : **41652952452**

Option 4 ID : **41652952450**

Status : **Answered**

Chosen Option : **4**

Q.26

A solid sphere, of radius R acquires a terminal velocity v_1 when falling (due to gravity) through a viscous fluid having a coefficient of viscosity η . The sphere is broken into 27 identical solid spheres. If each of these spheres acquires a terminal velocity, v_2 , when falling through the same fluid, the ratio (v_1/v_2) equals :

- Options
1. $1/9$
 2. 27
 3. $1/27$
 4. 9 ✓

Question Type : **MCQ**

Question ID : **41652913425**

Option 1 ID : **41652952479**

Option 2 ID : **41652952480**


Option 3 ID : **41652952481**

Option 4 ID : **41652952478**

Status : **Answered**

Chosen Option : **3**

Q.27 In the given circuit, the charge on $4 \mu\text{F}$ capacitor will be :



- Options
1. $9.6 \mu\text{C}$
 2. $5.4 \mu\text{C}$
 3. $24 \mu\text{C}$ ✓
 4. $13.4 \mu\text{C}$

Question Type : **MCQ**

Question ID : **41652913431**

Option 1 ID : **41652952502**

Option 2 ID : **41652952504**

Option 3 ID : **41652952503**

Option 4 ID : **41652952505**

Status : **Answered**

Chosen Option : **3**

Q.28 A system of three polarizers P_1, P_2, P_3 is set up such that the pass axis of P_3 is crossed with respect to that of P_1 . The pass axis of P_2 is inclined at 60° to the pass axis of P_3 . When a beam of unpolarized light of intensity I_0 is incident on P_1 , the intensity of light transmitted by the three polarizers is I . The ratio (I_0/I) equals (nearly) :

- Options
1. 5.33
 2. 16.00
 3. 1.80
 4. 10.67 ✓

Question Type : **MCQ**

Question ID : **41652913438**

Option 1 ID : **41652952531**

Option 2 ID : **41652952533**

Option 3 ID : **41652952530**

Option 4 ID : **41652952532**

Status : **Not Answered**

Chosen Option : --

Q.29 One kg of water, at 20°C , is heated in an electric kettle whose heating element has a mean (temperature averaged) resistance of 20Ω . The rms voltage in the mains is 200 V. Ignoring heat loss from the kettle, time taken for water to evaporate fully, is close to :

[Specific heat of water = $4200 \text{ J}/(\text{kg } ^\circ\text{C})$,
Latent heat of water = $2260 \text{ kJ}/\text{kg}$]

- Options
1. 3 minutes
 2. 16 minutes
 3. 22 minutes ✓
 4. 10 minutes

Question Type : **MCQ**

Question ID : **41652913432**

Option 1 ID : **41652952506**

Option 2 ID : **41652952509**

Option 3 ID : **41652952508**

Option 4 ID : **41652952507**

Status : **Not Answered**

Chosen Option : --

Q.30 A plane electromagnetic wave having a frequency $\nu = 23.9$ GHz propagates along the positive z -direction in free space. The peak value of the Electric Field is 60 V/m. Which among the following is the acceptable magnetic field component in the electromagnetic wave ?

- Options
1. $\vec{B} = 2 \times 10^{-7} \sin(0.5 \times 10^3 z - 1.5 \times 10^{11} t) \hat{i}$ ✓
 2. $\vec{B} = 2 \times 10^{-7} \sin(1.5 \times 10^2 x + 0.5 \times 10^{11} t) \hat{j}$
 3. $\vec{B} = 2 \times 10^7 \sin(0.5 \times 10^3 z + 1.5 \times 10^{11} t) \hat{i}$
 4. $\vec{B} = 60 \sin(0.5 \times 10^3 x + 1.5 \times 10^{11} t) \hat{k}$

Question Type : **MCQ**

Question ID : **41652913436**

Option 1 ID : **41652952523**

Option 2 ID : **41652952525**

Option 3 ID : **41652952524**





Option 4 ID : **41652952522**

Status : **Not Answered**

Chosen Option : --

Section : Chemistry

Q.1 Benzene diazonium chloride on reaction with aniline in the presence of dilute hydrochloric acid gives :

- Options
1.  ✓
 2. 
 3. 
 4. 

Question Type : **MCQ**

Question ID : **41652913451**

Option 1 ID : **41652952582**

Option 2 ID : **41652952584**

Option 3 ID : **41652952583**

Option 4 ID : 41652952585

Status : Answered

Chosen Option : 2

Q.2 A solution is prepared by dissolving 0.6 g of urea (molar mass = 60 g mol^{-1}) and 1.8 g of glucose (molar mass = 180 g mol^{-1}) in 100 mL of water at 27°C . The osmotic pressure of the solution is :
($R = 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$)

- Options
1. 8.2 atm
 2. 2.46 atm
 3. 4.92 atm ✓
 4. 1.64 atm

Question Type : MCQ

Question ID : 41652913470

Option 1 ID : 41652952661

Option 2 ID : 41652952659

Option 3 ID : 41652952658

Option 4 ID : 41652952660

Status : Answered

Chosen Option : 3

Q.3 The temporary hardness of a water sample is due to compound X. Boiling this sample converts X to compound Y. X and Y, respectively, are :

- Options
1. $\text{Mg}(\text{HCO}_3)_2$ and MgCO_3
 2. $\text{Ca}(\text{HCO}_3)_2$ and $\text{Ca}(\text{OH})_2$
 3. $\text{Mg}(\text{HCO}_3)_2$ and $\text{Mg}(\text{OH})_2$ ✓
 4. $\text{Ca}(\text{HCO}_3)_2$ and CaO

Question Type : MCQ

Question ID : 41652913458

Option 1 ID : 41652952611

Option 2 ID : 41652952612

Option 3 ID : 41652952610

Option 4 ID : 41652952613

Status : Answered

Chosen Option : 4


Q.4 The pair that has similar atomic radii is :

- Options
1. Mo and W ✓

2. Sc and Ni
3. Mn and Re
4. Ti and Hf

Question Type : **MCQ**Question ID : **41652913462**Option 1 ID : **41652952629**Option 2 ID : **41652952628**Option 3 ID : **41652952627**Option 4 ID : **41652952626**Status : **Answered**Chosen Option : **1**

Q.5 Consider the following reactions :



'A' is :

- Options
1. $\text{CH}_2=\text{CH}_2$
 2. $\text{CH}\equiv\text{CH}$
 3. $\text{CH}_3-\text{C}\equiv\text{C}-\text{CH}_3$
 4. $\text{CH}_3-\text{C}\equiv\text{CH}$ ✓

Question Type : **MCQ**Question ID : **41652913448**Option 1 ID : **41652952573**Option 2 ID : **41652952570**Option 3 ID : **41652952572**Option 4 ID : **41652952571**Status : **Not Answered**Chosen Option : **--**

Q.6 NO_2 required for a reaction is produced by the decomposition of N_2O_5 in CCl_4 as per the equation,



The initial concentration of N_2O_5 is 3.00 mol L^{-1} and it is 2.75 mol L^{-1} after 30 minutes. The rate of formation of NO_2 is :

- Options
1. $1.667 \times 10^{-2} \text{ mol L}^{-1} \text{ min}^{-1}$ ✓
 2. $8.333 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$

3. $4.167 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$

4. $2.083 \times 10^{-3} \text{ mol L}^{-1} \text{ min}^{-1}$

Question Type : **MCQ**Question ID : **41652913474**Option 1 ID : **41652952677**Option 2 ID : **41652952675**Option 3 ID : **41652952674**Option 4 ID : **41652952676**Status : **Answered**Chosen Option : **4****Q.7** The correct statement is :

- Options
1. pig iron is obtained from cast iron.
the Hall-Heroult process is used for
 2. the production of aluminium and iron.
leaching of bauxite using concentrated NaOH solution gives sodium aluminate and sodium silicate.
 3. ✓
 4. the blistered appearance of copper during the metallurgical process is due to the evolution of CO_2 .

Question Type : **MCQ**Question ID : **41652913457**Option 1 ID : **41652952607**Option 2 ID : **41652952609**Option 3 ID : **41652952608**Option 4 ID : **41652952606**Status : **Answered**Chosen Option : **2****Q.8** The C-C bond length is maximum in :

- Options
1. C_{70}
 2. C_{60}
 3. graphite
 4. diamond ✓

Question Type : **MCQ**Question ID : **41652913460**Option 1 ID : **41652952621**Option 2 ID : **41652952618**Option 3 ID : **41652952619**Option 4 ID : **41652952620**

Status : **Answered**
Chosen Option : 2

Q.9 The molar solubility of $\text{Cd}(\text{OH})_2$ is $1.84 \times 10^{-5} \text{ M}$ in water. The expected solubility of $\text{Cd}(\text{OH})_2$ in a buffer solution of $\text{pH} = 12$ is :

- Options
1. $2.49 \times 10^{-10} \text{ M}$ ✓
 2. $\frac{2.49}{1.84} \times 10^{-9} \text{ M}$
 3. $1.84 \times 10^{-9} \text{ M}$
 4. $6.23 \times 10^{-11} \text{ M}$

Question Type : **MCQ**
Question ID : **41652913472**
Option 1 ID : **41652952667**
Option 2 ID : **41652952668**
Option 3 ID : **41652952669**
Option 4 ID : **41652952666**
Status : **Not Answered**
Chosen Option : --

Q.10 The decreasing order of electrical conductivity of the following aqueous solutions is :

- 0.1 M Formic acid (A),
0.1 M Acetic acid (B),
0.1 M Benzoic acid (C).

- Options
1. $A > B > C$
 2. $A > C > B$ ✓
 3. $C > B > A$
 4. $C > A > B$

Question Type : **MCQ**
Question ID : **41652913473**
Option 1 ID : **41652952670**
Option 2 ID : **41652952673**
Option 3 ID : **41652952672**
Option 4 ID : **41652952671**
Status : **Answered**
Chosen Option : 3

Q.11

The coordination numbers of Co and Al in $[\text{Co}(\text{Cl})(\text{en})_2]\text{Cl}$ and $\text{K}_3[\text{Al}(\text{C}_2\text{O}_4)_3]$, respectively, are :

(en = ethane-1, 2-diamine)

- Options
1. 3 and 3
 2. 5 and 3
 3. 5 and 6 ✓
 4. 6 and 6

Question Type : MCQ

Question ID : 41652913464

Option 1 ID : 41652952635

Option 2 ID : 41652952637

Option 3 ID : 41652952636

Option 4 ID : 41652952634

Status : Answered

Chosen Option : 3

Q.12 Which of the given statements is INCORRECT about glycogen ?

- Options
1. It is present in animal cells.
 2. It is a straight chain polymer similar to amylose. ✓
 3. It is present in some yeast and fungi.
 4. Only α -linkages are present in the molecule.

Question Type : MCQ

Question ID : 41652913449

Option 1 ID : 41652952574

Option 2 ID : 41652952575

Option 3 ID : 41652952576

Option 4 ID : 41652952577

Status : Answered

Chosen Option : 2

Q.13 In which one of the following equilibria, $K_p \neq K_c$?

- Options
1. $2\text{HI}(\text{g}) \rightleftharpoons \text{H}_2(\text{g}) + \text{I}_2(\text{g})$
 2. $2\text{C}(\text{s}) + \text{O}_2(\text{g}) \rightleftharpoons 2\text{CO}(\text{g})$ ✓
 3. $2\text{NO}(\text{g}) \rightleftharpoons \text{N}_2(\text{g}) + \text{O}_2(\text{g})$
 4. $\text{NO}_2(\text{g}) + \text{SO}_2(\text{g}) \rightleftharpoons \text{NO}(\text{g}) + \text{SO}_3(\text{g})$

Question Type : **MCQ**Question ID : **41652913471**Option 1 ID : **41652952663**Option 2 ID : **41652952664**Option 3 ID : **41652952665**Option 4 ID : **41652952662**Status : **Answered**Chosen Option : **2****Q.14** In comparison to boron, beryllium has :

Options

1. greater nuclear charge and lesser first ionisation enthalpy.
2. lesser nuclear charge and greater first ionisation enthalpy. ✓
3. lesser nuclear charge and lesser first ionisation enthalpy.
4. greater nuclear charge and greater first ionisation enthalpy.

Question Type : **MCQ**Question ID : **41652913456**Option 1 ID : **41652952605**Option 2 ID : **41652952603**Option 3 ID : **41652952604**Option 4 ID : **41652952602**Status : **Answered**Chosen Option : **1****Q.15** The INCORRECT statement is :

Options

1. LiNO_3 decomposes on heating to give LiNO_2 and O_2 . ✓
2. LiCl crystallises from aqueous solution as $\text{LiCl} \cdot 2\text{H}_2\text{O}$.
3. Lithium is the strongest reducing agent among the alkali metals.
4. Lithium is least reactive with water among the alkali metals.

Question Type : **MCQ**Question ID : **41652913459**Option 1 ID : **41652952614**Option 2 ID : **41652952617**Option 3 ID : **41652952615**Option 4 ID : **41652952616**Status : **Answered**Chosen Option : **1**

Q.16 Thermal decomposition of a Mn compound (X) at 513 K results in compound Y, MnO_2 and a gaseous product. MnO_2 reacts with NaCl and concentrated H_2SO_4 to give a pungent gas Z. X, Y, and Z, respectively, are :

- Options
1. K_3MnO_4 , K_2MnO_4 and Cl_2
 2. K_2MnO_4 , KMnO_4 and Cl_2
 3. K_2MnO_4 , KMnO_4 and SO_2
 4. KMnO_4 , K_2MnO_4 and Cl_2 ✓

Question Type : MCQ

Question ID : 41652913461

Option 1 ID : 41652952625

Option 2 ID : 41652952624

Option 3 ID : 41652952622

Option 4 ID : 41652952623

Status : Not Answered

Chosen Option : --

Q.17 25 g of an unknown hydrocarbon upon burning produces 88 g of CO_2 and 9 g of H_2O . This unknown hydrocarbon contains :

- Options
1. 24 g of carbon and 1 g of hydrogen ✓
 2. 18 g of carbon and 7 g of hydrogen
 3. 22 g of carbon and 3 g of hydrogen
 4. 20 g of carbon and 5 g of hydrogen

Question Type : MCQ

Question ID : 41652913466

Option 1 ID : 41652952643

Option 2 ID : 41652952645

Option 3 ID : 41652952644

Option 4 ID : 41652952642

Status : Answered

Chosen Option : 1

Q.18 The INCORRECT match in the following is :

- Options
1. $\Delta G^0 < 0$, $K > 1$
 2. $\Delta G^0 > 0$, $K < 1$
 3. $\Delta G^0 = 0$, $K = 1$

4. $\Delta G^0 < 0, K < 1$ ✓

Question Type : **MCQ**

Question ID : **41652913469**

Option 1 ID : **41652952654**

Option 2 ID : **41652952656**

Option 3 ID : **41652952657**

Option 4 ID : **41652952655**

Status : **Answered**

Chosen Option : **4**

Q.19 An 'Assertion' and a 'Reason' are given below. Choose the correct answer from the following options :

Assertion (A): Vinyl halides do not undergo nucleophilic substitution easily.

Reason (R): Even though the intermediate carbocation is stabilized by loosely held π -electrons, the cleavage is difficult because of strong bonding.

Options

- Both (A) and (R) are correct statements but (R) is not the correct explanation of (A).
- Both (A) and (R) are wrong statements.
- Both (A) and (R) are correct statements and (R) is the correct explanation of (A).
- (A) is a correct statement but (R) is a wrong statement. ✓

Question Type : **MCQ**

Question ID : **41652913454**

Option 1 ID : **41652952595**

Option 2 ID : **41652952597**

Option 3 ID : **41652952594**

Option 4 ID : **41652952596**

Status : **Answered**

Chosen Option : **3**

Q.20 Among the following, the **INCORRECT** statement about colloids is :

Options

- They are larger than small molecules and have high molar mass.

2. They can scatter light.

The osmotic pressure of a colloidal solution is of higher order than the true solution at the same concentration. ✓

4. The range of diameters of colloidal particles is between 1 and 1000 nm.

Question Type : **MCQ**

Question ID : **41652913475**

Option 1 ID : **41652952681**

Option 2 ID : **41652952679**


Option 3 ID : **41652952680**

Option 4 ID : **41652952678**

Status : **Answered**

Chosen Option : **3**

- Q.21 The correct name of the following polymer is :



- Options
1. Polyisobutylene ✓
 2. Polytert-butylene
 3. Polyisobutane
 4. Polyisoprene

Question Type : **MCQ**

Question ID : **41652913450**

Option 1 ID : **41652952579**

Option 2 ID : **41652952580**

Option 3 ID : **41652952578**

Option 4 ID : **41652952581**

Status : **Answered**

Chosen Option : **2**

- Q.22 The ratio of number of atoms present in a simple cubic, body centered cubic and face centered cubic structure are, respectively :

- Options
1. 4 : 2 : 1
 2. 1 : 2 : 4 ✓
 3. 4 : 2 : 3
 4. 8 : 1 : 6





Question Type : **MCQ**Question ID : **41652913467**Option 1 ID : **41652952649**Option 2 ID : **41652952647**Option 3 ID : **41652952648**Option 4 ID : **41652952646**Status : **Answered**Chosen Option : **2**

Q.23 The primary pollutant that leads to photochemical smog is :

- Options
1. nitrogen oxides ✓
 2. ozone
 3. acrolein
 4. sulphur dioxide

Question Type : **MCQ**Question ID : **41652913465**Option 1 ID : **41652952641**Option 2 ID : **41652952638**Option 3 ID : **41652952640**Option 4 ID : **41652952639**Status : **Answered**Chosen Option : **1**

Q.24 Heating of 2-chloro-1-phenylbutane with EtOK/EtOH gives X as the major product. Reaction of X with $\text{Hg}(\text{OAc})_2/\text{H}_2\text{O}$ followed by NaBH_4 gives Y as the major product. Y is :

- Options
1.  ✓
 2. 
 3. 
 4. 

Question Type : **MCQ**Question ID : **41652913452**Option 1 ID : **41652952587**Option 2 ID : **41652952589**Option 3 ID : **41652952588**Option 4 ID : **41652952586**


Status : **Answered**
Chosen Option : **2**

Q.25 Among the following, the energy of 2s orbital is lowest in :

- Options
1. K ✓
 2. H
 3. Na
 4. Li

Question Type : **MCQ**
Question ID : **41652913468**
Option 1 ID : **41652952651**
Option 2 ID : **41652952650**
Option 3 ID : **41652952652**
Option 4 ID : **41652952653**
Status : **Answered**
Chosen Option : **2**

Q.26 The IUPAC name for the following compound is :



- Options
1. 3-methyl-4-(1-methylprop-2-ynyl)-1-heptene
 2. 3,5-dimethyl-4-propylhept-6-en-1-yne
 3. 3,5-dimethyl-4-propylhept-1-en-6-yne ✓
 4. 3-methyl-4-(3-methylprop-1-enyl)-1-heptyne

Question Type : **MCQ**
Question ID : **41652913447**
Option 1 ID : **41652952569**
Option 2 ID : **41652952566**
Option 3 ID : **41652952568**
Option 4 ID : **41652952567**
Status : **Answered**
Chosen Option : **3**

Q.27 Which one of the following is likely to give a precipitate with AgNO_3 solution ?

- Options
1. $\text{CH}_2 = \text{CH} - \text{Cl}$
 2. $(\text{CH}_3)_3\text{CCl}$ ✓
 3. CCl_4
 4. CHCl_3

Question Type : MCQ
Question ID : 41652913455
Option 1 ID : 41652952601
Option 2 ID : 41652952600
Option 3 ID : 41652952598
Option 4 ID : 41652952599
Status : Answered
Chosen Option : 4


Q.28 The compound used in the treatment of lead poisoning is :




- Options
1. desferrioxime B
 2. EDTA ✓
 3. Cis-platin
 4. D-penicillamine

Question Type : MCQ
Question ID : 41652913463
Option 1 ID : 41652952633
Option 2 ID : 41652952631
Option 3 ID : 41652952632
Option 4 ID : 41652952630
Status : Answered
Chosen Option : 3

Q.29 What will be the major product when m-cresol is reacted with propargyl bromide ($\text{HC} \equiv \text{C} - \text{CH}_2\text{Br}$) in presence of K_2CO_3 in acetone ?

Options





Question Type : **MCQ**

Question ID : **41652913453**

Option 1 ID : **41652952592**

Option 2 ID : **41652952590**


Option 3 ID : **41652952591**

Option 4 ID : **41652952593**

Status : **Answered**

Chosen Option : **3**

Q.30 In the following skew conformation of ethane, $H' - C - C - H''$ dihedral angle is :



Options 1. 58°

2. 120°

3. 149° ✓

4. 151°

Question Type : **MCQ**

Question ID : **41652913446**

Option 1 ID : **41652952562**

Option 2 ID : **41652952565**

Option 3 ID : **41652952564**

Option 4 ID : **41652952563**

Status : **Not Answered**

Chosen Option : --

Section : Mathematics

Q.1 Let $\alpha \in \mathbb{R}$ and the three vectors

$$\vec{a} = \alpha \hat{i} + \hat{j} + 3\hat{k}, \quad \vec{b} = 2\hat{i} + \hat{j} - \alpha\hat{k}$$

and $\vec{c} = \alpha\hat{i} - 2\hat{j} + 3\hat{k}$. Then the set

$$S = \{\alpha : \vec{a}, \vec{b} \text{ and } \vec{c} \text{ are coplanar}\}$$

- Options
1. is singleton
 2. contains exactly two positive numbers
 3. is empty ✓
 4. contains exactly two numbers only one of which is positive

Question Type : MCQ

Question ID : 41652913500

Option 1 ID : 41652952779

Option 2 ID : 41652952780

Option 3 ID : 41652952778

Option 4 ID : 41652952781

Status : Answered

Chosen Option : 3

Q.2 A person throws two fair dice. He wins Rs. 15 for throwing a doublet (same numbers on the two dice), wins Rs. 12 when the throw results in the sum of 9, and loses Rs. 6 for any other outcome on the throw. Then the expected gain/loss (in Rs.) of the person is :

- Options
1. $\frac{1}{2}$ loss ✓
 2. $\frac{1}{2}$ gain
 3. 2 gain
 4. $\frac{1}{4}$ loss

Question Type : MCQ

Question ID : 41652913501

Option 1 ID : 41652952783

Option 2 ID : 41652952784

Option 3 ID : 41652952782

Option 4 ID : 41652952785

Status : Not Answered

Chosen Option : --

Q.3 For an initial screening of an admission test, a candidate is given fifty problems to solve. If the probability that the candidate can solve any problem is $\frac{4}{5}$, then the probability that he is unable to solve less than two problems is :

Options

1. $\frac{201}{5} \left(\frac{1}{5}\right)^{49}$
2. $\frac{164}{25} \left(\frac{1}{5}\right)^{48}$
3. $\frac{316}{25} \left(\frac{4}{5}\right)^{48}$
4. $\frac{54}{5} \left(\frac{4}{5}\right)^{49}$ ✓

Question Type : MCQ

Question ID : 41652913502

Option 1 ID : 41652952786

Option 2 ID : 41652952789

Option 3 ID : 41652952788

Option 4 ID : 41652952787

Status : Not Answered

Chosen Option : --

Q.4 Let S be the set of all $\alpha \in \mathbb{R}$ such that the equation, $\cos 2x + \alpha \sin x = 2\alpha - 7$ has a solution. Then S is equal to :

Options

1. [3, 7]
2. [2, 6] ✓
3. [1, 4]
4. \mathbb{R}

Question Type : MCQ

Question ID : 41652913503

Option 1 ID : 41652952793

Option 2 ID : 41652952791

Option 3 ID : 41652952792

Option 4 ID : 41652952790

Status : **Answered**
Chosen Option : **2**

Q.5 If α , β and γ are three consecutive terms of a non-constant G.P. such that the equations $\alpha x^2 + 2\beta x + \gamma = 0$ and $x^2 + x - 1 = 0$ have a common root, then $\alpha(\beta + \gamma)$ is equal to :

- Options
1. $\beta\gamma$ ✓
 2. $\alpha\beta$
 3. $\alpha\gamma$
 4. 0

Question Type : **MCQ**
Question ID : **41652913478**
Option 1 ID : **41652952690**
Option 2 ID : **41652952691**
Option 3 ID : **41652952692**
Option 4 ID : **41652952693**
Status : **Not Answered**
Chosen Option : --

Q.6 A plane which bisects the angle between the two given planes $2x - y + 2z - 4 = 0$ and $x + 2y + 2z - 2 = 0$, passes through the point :

- Options
1. (2, 4, 1)
 2. (1, -4, 1)
 3. (1, 4, -1)
 4. (2, -4, 1) ✓

Question Type : **MCQ**
Question ID : **41652913498**
Option 1 ID : **41652952771**
Option 2 ID : **41652952770**
Option 3 ID : **41652952773**
Option 4 ID : **41652952772**
Status : **Not Answered**
Chosen Option : --

Q.7 Let $\alpha \in (0, \pi/2)$ be fixed. If the integral

$$\int \frac{\tan x + \tan \alpha}{\tan x - \tan \alpha} dx =$$

$A(x) \cos 2\alpha + B(x) \sin 2\alpha + C$, where C is a constant of integration, then the functions $A(x)$ and $B(x)$ are respectively :

Options

1. $x - \alpha$ and $\log_e |\sin(x - \alpha)|$ ✓
2. $x + \alpha$ and $\log_e |\sin(x - \alpha)|$
3. $x + \alpha$ and $\log_e |\sin(x + \alpha)|$
4. $x - \alpha$ and $\log_e |\cos(x - \alpha)|$

Question Type : MCQ

Question ID : 41652913489

Option 1 ID : 41652952734

Option 2 ID : 41652952735

Option 3 ID : 41652952737

Option 4 ID : 41652952736

Status : Answered

Chosen Option : 2

Q.8

$$\lim_{x \rightarrow 0} \frac{x + 2 \sin x}{\sqrt{x^2 + 2 \sin x + 1} - \sqrt{\sin^2 x - x + 1}}$$

is :

Options

1. 3
2. 1
3. 2 ✓
4. 6

Question Type : MCQ

Question ID : 41652913485

Option 1 ID : 41652952720

Option 2 ID : 41652952718

Option 3 ID : 41652952719

Option 4 ID : 41652952721

Status : Answered

Chosen Option : 3

Q.9

A group of students comprises of 5 boys and n girls. If the number of ways, in which a team of 3 students can randomly be selected from this group such that there is at least one boy and at least one girl in each team, is 1750, then n is equal to :

Options

1. 24
2. 27
3. 25 ✓
4. 28

Question Type : MCQ

Question ID : **41652913481**
 Option 1 ID : **41652952702**
 Option 2 ID : **41652952704**
 Option 3 ID : **41652952703**
 Option 4 ID : **41652952705**
 Status : **Answered**
 Chosen Option : **3**

Q.10 The equation of a common tangent to the curves, $y^2 = 16x$ and $xy = -4$, is :

- Options
1. $x - 2y + 16 = 0$
 2. $x - y + 4 = 0$ ✓
 3. $2x - y + 2 = 0$
 4. $x + y + 4 = 0$

Question Type : **MCQ**
 Question ID : **41652913496**
 Option 1 ID : **41652952762**
 Option 2 ID : **41652952765**
 Option 3 ID : **41652952764**
 Option 4 ID : **41652952763**
 Status : **Not Answered**
 Chosen Option : **--**

Q.11 A circle touching the x -axis at $(3, 0)$ and making an intercept of length 8 on the y -axis passes through the point :

- Options
1. $(3, 10)$ ✓
 2. $(2, 3)$
 3. $(3, 5)$
 4. $(1, 5)$

Question Type : **MCQ**
 Question ID : **41652913495**
 Option 1 ID : **41652952758**
 Option 2 ID : **41652952761**
 Option 3 ID : **41652952759**
 Option 4 ID : **41652952760**
 Status : **Answered**
 Chosen Option : **1**

Q.12 Let A, B and C be sets such that $\phi \neq A \cap B \subseteq C$. Then which of the following statements is not true ?

- Options
1. $B \cap C \neq \phi$
 2. $(C \cup A) \cap (C \cup B) = C$

3. If $(A - B) \subseteq C$, then $A \subseteq C$
4. If $(A - C) \subseteq B$, then $A \subseteq B$ ✓

Question Type : MCQ

Question ID : 41652913476

Option 1 ID : 41652952682

Option 2 ID : 41652952683

Option 3 ID : 41652952684

Option 4 ID : 41652952685

Status : Answered

Chosen Option : 4

- Q.13 Let $f(x) = 5 - |x - 2|$ and $g(x) = |x + 1|$,
 $x \in \mathbb{R}$. If $f(x)$ attains maximum value at α
 and $g(x)$ attains minimum value at β , then

$$\lim_{x \rightarrow -\alpha\beta} \frac{(x-1)(x^2 - 5x + 6)}{x^2 - 6x + 8} \text{ is equal to :}$$

- Options
1. $3/2$
 2. $1/2$ ✓
 3. $-3/2$
 4. $-1/2$

Question Type : MCQ

Question ID : 41652913488

Option 1 ID : 41652952732

Option 2 ID : 41652952730

Option 3 ID : 41652952733

Option 4 ID : 41652952731

Status : Answered

Chosen Option : 2

- Q.14 Let $z \in \mathbb{C}$ with $\text{Im}(z) = 10$ and it satisfies

$$\frac{2z - n}{2z + n} = 2i - 1 \text{ for some natural number}$$

n. Then :

- Options
1. $n = 20$ and $\text{Re}(z) = 10$
 2. $n = 40$ and $\text{Re}(z) = 10$
 3. $n = 20$ and $\text{Re}(z) = -10$
 4. $n = 40$ and $\text{Re}(z) = -10$ ✓

Question Type : MCQ

Question ID : 41652913477

Option 1 ID : 41652952686

Option 2 ID : 41652952688

Option 3 ID : 41652952687

Option 4 ID : 41652952689

Status : Answered

Chosen Option : 4

Q.15 The tangents to the curve $y = (x - 2)^2 - 1$ at its points of intersection with the line $x - y = 3$, intersect at the point :

Options

1. $\left(\frac{5}{2}, 1\right)$
2. $\left(\frac{5}{2}, -1\right)$ ✓
3. $\left(-\frac{5}{2}, -1\right)$
4. $\left(-\frac{5}{2}, 1\right)$

Question Type : MCQ

Question ID : 41652913487

Option 1 ID : 41652952728

Option 2 ID : 41652952726

Option 3 ID : 41652952727

Option 4 ID : 41652952729

Status : Answered

Chosen Option : 2

Q.16 The derivative of $\tan^{-1}\left(\frac{\sin x - \cos x}{\sin x + \cos x}\right)$, with respect to $\frac{x}{2}$, where $\left(x \in \left(0, \frac{\pi}{2}\right)\right)$ is:

Options

1. 2 ✓
2. $\frac{1}{2}$
3. $\frac{2}{3}$
4. 1

Question Type : MCQ

Question ID : 41652913486

Option 1 ID : 41652952724

Option 2 ID : 41652952723

Option 3 ID : 41652952725

Option 4 ID : 41652952722

Status : Not Answered

Chosen Option : --

Q.17

If a_1, a_2, a_3, \dots are in A.P. such that $a_1 + a_7 + a_{16} = 40$, then the sum of the first 15 terms of this A.P. is :

- Options
1. 280
 2. 120
 3. 150
 4. 200 ✓

Question Type : **MCQ**

Question ID : **41652913483**

Option 1 ID : **41652952712**

Option 2 ID : **41652952711**

Option 3 ID : **41652952710**

Option 4 ID : **41652952713**

Status : **Answered**

Chosen Option : **4**

Q.18 If ${}^{20}C_1 + (2^2) {}^{20}C_2 + (3^2) {}^{20}C_3 + \dots + (20^2) {}^{20}C_{20} = A(2^\beta)$, then the ordered pair (A, β) is equal to :

- Options
1. (380, 19)
 2. (420, 18) ✓
 3. (420, 19)
 4. (380, 18)

Question Type : **MCQ**

Question ID : **41652913484**

Option 1 ID : **41652952715**

Option 2 ID : **41652952716**

Option 3 ID : **41652952717**

Option 4 ID : **41652952714**

Status : **Answered**

Chosen Option : **3**

Q.19 The term independent of x in the expansion

of $\left(\frac{1}{60} - \frac{x^8}{81}\right) \cdot \left(2x^2 - \frac{3}{x^2}\right)^6$ is equal to :

- Options
1. -72
 2. 36
 3. -108
 4. -36 ✓

Question Type : **MCQ**

Question ID : **41652913482**
 Option 1 ID : **41652952707**
 Option 2 ID : **41652952709**
 Option 3 ID : **41652952706**
 Option 4 ID : **41652952708**
 Status : **Not Answered**
 Chosen Option : --

Q.20 If the area (in sq. units) bounded by the parabola $y^2 = 4\lambda x$ and the line $y = \lambda x$, $\lambda > 0$, is $\frac{1}{9}$, then λ is equal to :

- Options
1. $4\sqrt{3}$
 2. $2\sqrt{6}$
 3. 48
 4. 24 ✓

Question Type : **MCQ**
 Question ID : **41652913491**
 Option 1 ID : **41652952742**
 Option 2 ID : **41652952743**
 Option 3 ID : **41652952745**
 Option 4 ID : **41652952744**
 Status : **Answered**
 Chosen Option : 4

Q.21 The general solution of the differential equation $(y^2 - x^3) dx - xy dy = 0$ ($x \neq 0$) is :
 (where c is a constant of integration)

- Options
1. $y^2 + 2x^2 + cx^3 = 0$
 2. $y^2 - 2x^2 + cx^3 = 0$
 3. $y^2 - 2x^3 + cx^2 = 0$
 4. $y^2 + 2x^3 + cx^2 = 0$ ✓

Question Type : **MCQ**
 Question ID : **41652913492**
 Option 1 ID : **41652952746**
 Option 2 ID : **41652952749**
 Option 3 ID : **41652952747**
 Option 4 ID : **41652952748**
 Status : **Not Answered**
 Chosen Option : --

Q.22

A triangle has a vertex at $(1, 2)$ and the midpoints of the two sides through it are $(-1, 1)$ and $(2, 3)$. Then the centroid of this triangle is :

Options

1. $\left(\frac{1}{3}, 1\right)$
2. $\left(1, \frac{7}{3}\right)$
3. $\left(\frac{1}{3}, 2\right)$ ✓
4. $\left(\frac{1}{3}, \frac{5}{3}\right)$

Question Type : **MCQ**

Question ID : **41652913494**

Option 1 ID : **41652952754**

Option 2 ID : **41652952756**

Option 3 ID : **41652952755**

Option 4 ID : **41652952757**

Status : **Answered**

Chosen Option : **3**

Q.23 The angle of elevation of the top of a vertical tower standing on a horizontal plane is observed to be 45° from a point A on the plane. Let B be the point 30 m vertically above the point A. If the angle of elevation of the top of the tower from B be 30° , then the distance (in m) of the foot of the tower from the point A is :

Options

1. $15(3 + \sqrt{3})$ ✓
2. $15(1 + \sqrt{3})$
3. $15(5 - \sqrt{3})$
4. $15(3 - \sqrt{3})$

Question Type : **MCQ**

Question ID : **41652913504**

Option 1 ID : **41652952795**

Option 2 ID : **41652952796**

Option 3 ID : **41652952797**

Option 4 ID : **41652952794**

Status : **Answered**

Chosen Option : **1**

Q.24 The Boolean expression $\sim(p \Rightarrow (\sim q))$ is equivalent to :

- Options
1. $(\sim p) \Rightarrow q$
 2. $q \Rightarrow \sim p$
 3. $p \vee q$
 4. $p \wedge q$ ✓

Question Type : MCQ

Question ID : 41652913505

Option 1 ID : 41652952798

Option 2 ID : 41652952799

Option 3 ID : 41652952801

Option 4 ID : 41652952800

Status : Answered

Chosen Option : 2

Q.25 The length of the perpendicular drawn from the point $(2, 1, 4)$ to the plane containing the lines

$$\vec{r} = (\hat{i} + \hat{j}) + \lambda(\hat{i} + 2\hat{j} - \hat{k}) \quad \text{and}$$

$$\vec{r} = (\hat{i} + \hat{j}) + \mu(-\hat{i} + \hat{j} - 2\hat{k}) \quad \text{is :}$$

- Options
1. $\frac{1}{3}$
 2. 3
 3. $\sqrt{3}$ ✓
 4. $\frac{1}{\sqrt{3}}$

Question Type : MCQ

Question ID : 41652913499

Option 1 ID : 41652952776

Option 2 ID : 41652952774

Option 3 ID : 41652952775

Option 4 ID : 41652952777

Status : Not Answered

Chosen Option : --

Q.26 A value of $\theta \in (0, \pi/3)$, for which

$$\begin{vmatrix} 1 + \cos^2 \theta & \sin^2 \theta & 4 \cos 6\theta \\ \cos^2 \theta & 1 + \sin^2 \theta & 4 \cos 6\theta \\ \cos^2 \theta & \sin^2 \theta & 1 + 4 \cos 6\theta \end{vmatrix} = 0, \text{ is :}$$

- Options
1. $\frac{\pi}{9}$ ✓
 2. $\frac{7\pi}{24}$
 3. $\frac{7\pi}{36}$
 4. $\frac{\pi}{18}$

Question Type : **MCQ**

Question ID : **41652913479**

Option 1 ID : **41652952695**

Option 2 ID : **41652952696**

Option 3 ID : **41652952697**

Option 4 ID : **41652952694**

Status : **Not Answered**

Chosen Option : --

Q.27 A straight line L at a distance of 4 units from the origin makes positive intercepts on the coordinate axes and the perpendicular from the origin to this line makes an angle of 60° with the line $x + y = 0$. Then an equation of the line L is :

- Options
1. $(\sqrt{3} + 1)x + (\sqrt{3} - 1)y = 8\sqrt{2}$ ✓
 2. $x + \sqrt{3}y = 8$
 3. $\sqrt{3}x + y = 8$
 4. $(\sqrt{3} - 1)x + (\sqrt{3} + 1)y = 8\sqrt{2}$

Question Type : **MCQ**

Question ID : **41652913493**

Option 1 ID : **41652952750**

Option 2 ID : **41652952753**

Option 3 ID : **41652952752**

Option 4 ID : **41652952751**

Status : **Not Answered**

Chosen Option : --

Q.28 An ellipse, with foci at $(0, 2)$ and $(0, -2)$ and minor axis of length 4, passes through which of the following points ?

- Options
1. $(1, 2\sqrt{2})$
 2. $(2, \sqrt{2})$

3. $(\sqrt{2}, 2)$ ✓

4. $(2, 2\sqrt{2})$

Question Type : **MCQ**Question ID : **41652913497**Option 1 ID : **41652952767**Option 2 ID : **41652952766**Option 3 ID : **41652952768**Option 4 ID : **41652952769**Status : **Answered**Chosen Option : **4**

Q.29

A value of α such that

$$\int_{\alpha}^{\alpha+1} \frac{dx}{(x+\alpha)(x+\alpha+1)} = \log_e \left(\frac{9}{8} \right) \text{ is :}$$

Options

1. $-\frac{1}{2}$

2. $\frac{1}{2}$

3. -2 ✓

4. 2

Question Type : **MCQ**Question ID : **41652913490**Option 1 ID : **41652952741**Option 2 ID : **41652952738**Option 3 ID : **41652952740**Option 4 ID : **41652952739**Status : **Answered**Chosen Option : **2**

Q.30

If $[x]$ denotes the greatest integer $\leq x$, then the system of linear equations

$$[\sin\theta]x + [-\cos\theta]y = 0$$

$$[\cot\theta]x + y = 0$$

Options

has a unique solution if

1. $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3} \right) \cup \left(\pi, \frac{7\pi}{6} \right)$.

have infinitely many solutions if

2. $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3} \right) \cup \left(\pi, \frac{7\pi}{6} \right)$.

has a unique solution if $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right)$

3. and have infinitely many solutions if

$$\theta \in \left(\pi, \frac{7\pi}{6}\right).$$

have infinitely many solutions if

4. $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right)$ and has a unique

solution if $\theta \in \left(\pi, \frac{7\pi}{6}\right)$. ✓

Question Type : **MCQ**

Question ID : **41652913480**

Option 1 ID : **41652952698**

Option 2 ID : **41652952699**

Option 3 ID : **41652952700**

Option 4 ID : **41652952701**

Status : **Answered**

Chosen Option : **2**