

JEE MAIN 2019


Application No	
Candidate Name	
Roll No.	
Test Date	10/01/2019
Test Time	2:30 PM - 5:30 PM
Subject	Paper I EH

Section : Physics

Q.1

The Wheatstone bridge shown in Fig. here, gets balanced when the carbon resistor used as R_1 has the colour code (Orange, Red, Brown). The resistors R_2 and R_4 are 80Ω and 40Ω , respectively.

Assuming that the colour code for the carbon resistors gives their accurate values, the colour code for the carbon resistor, used as R_3 , would be :



- Options
1. Brown, Blue, Brown
 2. Brown, Blue, Black
 3. Red, Green, Brown
 4. Grey, Black, Brown

Question ID : 41652910434

Option 1 ID : 41652941197

Option 2 ID : 41652941196


Option 3 ID : 41652941195

Option 4 ID : 41652941194

Status : Answered

Chosen Option : 2

Q.2 Consider the nuclear fission



Given that the binding energy/nucleon of Ne^{20} , He^4 and C^{12} are, respectively, 8.03 MeV , 7.07 MeV and 7.86 MeV , identify the correct statement :

- Options
1. energy of 12.4 MeV will be supplied
 2. 8.3 MeV energy will be released
 3. energy of 3.6 MeV will be released

4. energy of 11.9 MeV has to be supplied

Question ID : 41652910431

Option 1 ID : 41652941184

Option 2 ID : 41652941185

Option 3 ID : 41652941183

Option 4 ID : 41652941182

Status : Not Answered

Chosen Option : --

Q.3

A hoop and a solid cylinder of same mass and radius are made of a permanent magnetic material with their magnetic moment parallel to their respective axes. But the magnetic moment of hoop is twice of solid cylinder. They are placed in a uniform magnetic field in such a manner that their magnetic moments make a small angle with the field. If the oscillation periods of hoop and cylinder are T_h and T_c respectively, then :

- Options
1. $T_h = T_c$
 2. $T_h = 2T_c$
 3. $T_h = 1.5T_c$
 4. $T_h = 0.5T_c$

Question ID : 41652910424

Option 1 ID : 41652941156

Option 2 ID : 41652941155

Option 3 ID : 41652941157

Option 4 ID : 41652941154

Status : Not Answered

Chosen Option : --

Q.4

An unknown metal of mass 192 g heated to a temperature of 100°C was immersed into a brass calorimeter of mass 128 g containing 240 g of water at a temperature of 8.4°C . Calculate the specific heat of the unknown metal if water temperature stabilizes at 21.5°C . (Specific heat of brass is $394 \text{ J kg}^{-1} \text{ K}^{-1}$)

- Options
1. $458 \text{ J kg}^{-1} \text{ K}^{-1}$
 2. $1232 \text{ J kg}^{-1} \text{ K}^{-1}$
 3. $916 \text{ J kg}^{-1} \text{ K}^{-1}$
 4. $654 \text{ J kg}^{-1} \text{ K}^{-1}$

Question ID : 41652910435

Option 1 ID : 41652941199

Option 2 ID : 41652941201

Option 3 ID : 41652941198


Option 4 ID : 41652941200

Status : Not Answered

Chosen Option : --

Q.5

Two identical spherical balls of mass M and radius R each are stuck on two ends of a rod of length $2R$ and mass M (see figure). The moment of inertia of the system about the axis passing perpendicularly through the centre of the rod is :



Options

1. $\frac{137}{15} MR^2$
2. $\frac{17}{15} MR^2$
3. $\frac{209}{15} MR^2$
4. $\frac{152}{15} MR^2$

Question ID : 41652910412

Option 1 ID : 41652941107

Option 2 ID : 41652941106

Option 3 ID : 41652941109

Option 4 ID : 41652941108

Status : Answered

Chosen Option : 1

Q.6

The self induced emf of a coil is 25 volts. When the current in it is changed at uniform rate from 10 A to 25 A in 1 s, the change in the energy of the inductance is :

Options

1. 740 J
2. 437.5 J
3. 540 J
4. 637.5 J

Question ID : 41652910426

Option 1 ID : 41652941165

Option 2 ID : 41652941162

Option 3 ID : 41652941163

Option 4 ID : 41652941164


Status : Answered

Chosen Option : 3

Q.7

The actual value of resistance R , shown in the figure is $30\ \Omega$. This is measured in an experiment as shown using the standard

formula $R = \frac{V}{I}$, where V and I are the readings of the voltmeter and ammeter, respectively. If the measured value of R is 5% less, then the internal resistance of the voltmeter is :



- Options
1. $600\ \Omega$
 2. $570\ \Omega$
 3. $35\ \Omega$
 4. $350\ \Omega$

Question ID : 41652910423

Option 1 ID : 41652941153

Option 2 ID : 41652941151

Option 3 ID : 41652941150

Option 4 ID : 41652941152

Status : Answered

Chosen Option : 2

Q.8 At some location on earth the horizontal component of earth's magnetic field is $18 \times 10^{-6}\ \text{T}$. At this location, magnetic needle of length $0.12\ \text{m}$ and pole strength $1.8\ \text{Am}$ is suspended from its mid-point using a thread, it makes 45° angle with horizontal in equilibrium. To keep this needle horizontal, the vertical force that should be applied at one of its ends is :

- Options
1. $3.6 \times 10^{-5}\ \text{N}$
 2. $1.8 \times 10^{-5}\ \text{N}$
 3. $1.3 \times 10^{-5}\ \text{N}$
 4. $6.5 \times 10^{-5}\ \text{N}$

Question ID : 41652910425

Option 1 ID : 41652941161

Option 2 ID : 41652941159

Option 3 ID : 41652941160

Option 4 ID : 41652941158

Status : Answered

Chosen Option : 4

Q.9

Two vectors \vec{A} and \vec{B} have equal magnitudes. The magnitude of $(\vec{A} + \vec{B})$ is 'n' times the magnitude of $(\vec{A} - \vec{B})$. The angle between \vec{A} and \vec{B} is :

Options

1. $\cos^{-1} \left[\frac{n^2 - 1}{n^2 + 1} \right]$

2. $\cos^{-1} \left[\frac{n - 1}{n + 1} \right]$

3. $\sin^{-1} \left[\frac{n^2 - 1}{n^2 + 1} \right]$

4. $\sin^{-1} \left[\frac{n - 1}{n + 1} \right]$

Question ID : 41652910407

Option 1 ID : 41652941086

Option 2 ID : 41652941087

Option 3 ID : 41652941088

Option 4 ID : 41652941089

Status : Not Answered

Chosen Option : --

Q.10 A metal plate of area $1 \times 10^{-4} \text{ m}^2$ is illuminated by a radiation of intensity 16 mW/m^2 . The work function of the metal is 5 eV . The energy of the incident photons is 10 eV and only 10% of it produces photo electrons. The number of emitted photo electrons per second and their maximum energy, respectively, will be : [1 eV = $1.6 \times 10^{-19} \text{ J}$]

Options 1. 10^{14} and 10 eV 2. 10^{12} and 5 eV 3. 10^{11} and 5 eV 4. 10^{10} and 5 eV

Question ID : 41652910430

Option 1 ID : 41652941181

Option 2 ID : 41652941180

Option 3 ID : 41652941179

Option 4 ID : 41652941178

Status : Answered

Chosen Option : 3


Q.11

A particle which is experiencing a force, given by $\vec{F} = 3\vec{i} - 12\vec{j}$, undergoes a displacement of $\vec{d} = 4\vec{i}$. If the particle had a kinetic energy of 3 J at the beginning of the displacement, what is its kinetic energy at the end of the displacement?

- Options
1. 9 J
 2. 12 J
 3. 10 J
 4. 15 J

Question ID : 41652910410
 Option 1 ID : 41652941100
 Option 2 ID : 41652941099
 Option 3 ID : 41652941101
 Option 4 ID : 41652941098
 Status : Not Answered
 Chosen Option : --

Q.12 Consider a Young's double slit experiment as shown in figure. What should be the slit separation d in terms of wavelength λ such that the first minima occurs directly in front of the slit (S_1)?



- Options
1. $\frac{\lambda}{2(\sqrt{5} - 2)}$
 2. $\frac{\lambda}{(\sqrt{5} - 2)}$
 3. $\frac{\lambda}{2(5 - \sqrt{2})}$
 4. $\frac{\lambda}{(5 - \sqrt{2})}$

Question ID : 41652910429
 Option 1 ID : 41652941175
 Option 2 ID : 41652941174
 Option 3 ID : 41652941177
 Option 4 ID : 41652941176
 Status : Answered
 Chosen Option : 1

Q.13

The eye can be regarded as a single refracting surface. The radius of curvature of this surface is equal to that of cornea (7.8 mm). This surface separates two media of refractive indices 1 and 1.34. Calculate the distance from the refracting surface at which a parallel beam of light will come to focus.

- Options
- 1 cm
 - 2 cm
 - 4.0 cm
 - 3.1 cm

Question ID : 41652910428

Option 1 ID : 41652941171

Option 2 ID : 41652941170

Option 3 ID : 41652941173

Option 4 ID : 41652941172

Status : Not Answered

Chosen Option : --

Q.14 A current of 2 mA was passed through an unknown resistor which dissipated a power of 4.4 W. Dissipated power when an ideal power supply of 11 V is connected across it is :

- Options
1. 11×10^{-5} W
 2. 11×10^{-3} W
 3. 11×10^{-4} W
 4. 11×10^5 W

Question ID : 41652910422

Option 1 ID : 41652941148

Option 2 ID : 41652941146

Option 3 ID : 41652941147

Option 4 ID : 41652941149

Status : Not Answered

Chosen Option : --

Q.15 The diameter and height of a cylinder are measured by a meter scale to be 12.6 ± 0.1 cm and 34.2 ± 0.1 cm, respectively. What will be the value of its volume in appropriate significant figures ?

- Options
1. 4264 ± 81 cm³
 2. 4264.4 ± 81.0 cm³
 3. 4260 ± 80 cm³
 4. 4300 ± 80 cm³

Question ID : 41652910406

Option 1 ID : 41652941084

Option 2 ID : 41652941082

Option 3 ID : 41652941083

Option 4 ID : 41652941085
 Status : Not Answered
 Chosen Option : --

Q.16 Four equal point charges Q each are placed in the xy plane at $(0, 2)$, $(4, 2)$, $(4, -2)$ and $(0, -2)$. The work required to put a fifth charge Q at the origin of the coordinate system will be :

Options

1. $\frac{Q^2}{4\pi\epsilon_0} \left(1 + \frac{1}{\sqrt{3}}\right)$

2. $\frac{Q^2}{4\pi\epsilon_0} \left(1 + \frac{1}{\sqrt{5}}\right)$

3. $\frac{Q^2}{2\sqrt{2}\pi\epsilon_0}$

4. $\frac{Q^2}{4\pi\epsilon_0}$

Question ID : 41652910421
 Option 1 ID : 41652941145
 Option 2 ID : 41652941142
 Option 3 ID : 41652941143
 Option 4 ID : 41652941144
 Status : Not Answered
 Chosen Option : --

Q.17 The modulation frequency of an AM radio station is 250 kHz, which is 10% of the carrier wave. If another AM station approaches you for license what broadcast frequency will you allot ?

Options

1. 2750 kHz

2. 2900 kHz

3. 2250 kHz

4. 2000 kHz

Question ID : 41652910433
 Option 1 ID : 41652941192
 Option 2 ID : 41652941193
 Option 3 ID : 41652941191
 Option 4 ID : 41652941190
 Status : Not Answered
 Chosen Option : --


Q.18 A closed organ pipe has a fundamental frequency of 1.5 kHz. The number of overtones that can be distinctly heard by a person with this organ pipe will be : (Assume that the highest frequency a person can hear is 20,000 Hz)

Options 1. 6

2. 4
3. 7
4. 5

Question ID : 41652910418
 Option 1 ID : 41652941130
 Option 2 ID : 41652941132
 Option 3 ID : 41652941131
 Option 4 ID : 41652941133
 Status : Answered
 Chosen Option : 1

Q.19 For the circuit shown below, the current through the Zener diode is :



- Options
1. 9 mA
 2. 5 mA
 3. Zero
 4. 14 mA

Question ID : 41652910432
 Option 1 ID : 41652941189
 Option 2 ID : 41652941187
 Option 3 ID : 41652941186
 Option 4 ID : 41652941188
 Status : Not Answered
 Chosen Option : --

Q.20 The electric field of a plane polarized electromagnetic wave in free space at time $t=0$ is given by an expression

$$\vec{E}(x, y) = 10 \hat{j} \cos [(6x + 8z)]$$

The magnetic field $\vec{B}(x, z, t)$ is given by : (c is the velocity of light)

- Options
1. $\frac{1}{c} (6\hat{k} + 8\hat{i}) \cos [(6x - 8z + 10ct)]$
 2. $\frac{1}{c} (6\hat{k} - 8\hat{i}) \cos [(6x + 8z - 10ct)]$
 3. $\frac{1}{c} (6\hat{k} + 8\hat{i}) \cos [(6x + 8z - 10ct)]$
 4. $\frac{1}{c} (6\hat{k} - 8\hat{i}) \cos [(6x + 8z + 10ct)]$

Question ID : 41652910427

Option 1 ID : 41652941167

Option 2 ID : 41652941166

Option 3 ID : 41652941168

Option 4 ID : 41652941169


Status : Answered

Chosen Option : 4

Q.21 Charges $-q$ and $+q$ located at A and B, respectively, constitute an electric dipole. Distance $AB=2a$, O is the mid point of the dipole and OP is perpendicular to AB. A charge Q is placed at P where $OP=y$ and $y \gg 2a$. The charge Q experiences an electrostatic force F. If Q is now moved along the equatorial line to P' such that

$OP' = \left(\frac{y}{3}\right)$, the force on Q will be close

to: $\left(\frac{y}{3} \gg 2a\right)$

Options 1. $3F$ 2. $\frac{F}{3}$ 3. $9F$ 4. $27F$

Question ID : 41652910420

Option 1 ID : 41652941139

Option 2 ID : 41652941138

Option 3 ID : 41652941140

Option 4 ID : 41652941141

Status : Answered

Chosen Option : 3

Q.22 Two stars of masses 3×10^{31} kg each, and at distance 2×10^{11} m rotate in a plane about their common centre of mass O. A meteorite passes through O moving perpendicular to the star's rotation plane. In order to escape from the gravitational field of this double star, the minimum speed that meteorite should have at O is :

(Take Gravitational constant $G = 6.67 \times 10^{-11} \text{ Nm}^2 \text{ kg}^{-2}$)

Options

1. 2.4×10^4 m/s
2. 1.4×10^5 m/s
3. 3.8×10^4 m/s
4. 2.8×10^5 m/s

Question ID : 41652910413
Option 1 ID : 41652941113
Option 2 ID : 41652941112
Option 3 ID : 41652941111
Option 4 ID : 41652941110
Status : Answered
Chosen Option : 2

Q.23 Half mole of an ideal monoatomic gas is heated at constant pressure of 1 atm from 20°C to 90°C . Work done by gas is close to : (Gas constant $R = 8.31$ J/mol·K)

- Options
1. 581 J
 2. 291 J
 3. 146 J
 4. 73 J

Question ID : 41652910415
Option 1 ID : 41652941118
Option 2 ID : 41652941119
Option 3 ID : 41652941120
Option 4 ID : 41652941121
Status : Answered
Chosen Option : 2

Q.24 A parallel plate capacitor having capacitance 12 pF is charged by a battery to a potential difference of 10 V between its plates. The charging battery is now disconnected and a porcelain slab of dielectric constant 6.5 is slipped between the plates. The work done by the capacitor on the slab is :

- Options
1. 692 pJ
 2. 508 pJ
 3. 560 pJ
 4. 600 pJ

Question ID : 41652910419
Option 1 ID : 41652941136
Option 2 ID : 41652941134
Option 3 ID : 41652941137
Option 4 ID : 41652941135
Status : Answered
Chosen Option : 3

Q.25

A particle starts from the origin at time $t = 0$ and moves along the positive x -axis. The graph of velocity with respect to time is shown in figure. What is the position of the particle at time $t = 5\text{s}$?



- Options
1. 10 m
 2. 6 m
 3. 3 m
 4. 9 m

Question ID : 41652910408

Option 1 ID : 41652941092

Option 2 ID : 41652941091


Option 3 ID : 41652941090

Option 4 ID : 41652941093

Status : Answered

Chosen Option : 2

- Q.26 A rigid massless rod of length $3l$ has two masses attached at each end as shown in the figure. The rod is pivoted at point P on the horizontal axis (see figure). When released from initial horizontal position, its instantaneous angular acceleration will be :



- Options
1. $\frac{g}{13l}$
 2. $\frac{g}{3l}$
 3. $\frac{g}{2l}$
 4. $\frac{7g}{3l}$

Question ID : 41652910411

Option 1 ID : 41652941105

Option 2 ID : 41652941103

Option 3 ID : 41652941102

Option 4 ID : 41652941104

Status : Answered

Chosen Option : 2

Q.27 Two forces P and Q, of magnitude $2F$ and $3F$, respectively, are at an angle θ with each other. If the force Q is doubled, then their resultant also gets doubled. Then, the angle θ is :

- Options
1. 120°
 2. 60°
 3. 90°
 4. 30°

Question ID : 41652910409

Option 1 ID : 41652941097

Option 2 ID : 41652941095

Option 3 ID : 41652941096

Option 4 ID : 41652941094

Status : Answered

Chosen Option : 4

Q.28 A cylindrical plastic bottle of negligible mass is filled with 310 ml of water and left floating in a pond with still water. If pressed downward slightly and released, it starts performing simple harmonic motion at angular frequency ω . If the radius of the bottle is 2.5 cm then ω is close to : (density of water = 10^3 kg/m^3)

- Options
1. 3.75 rad s^{-1}
 2. 1.25 rad s^{-1}
 3. 2.50 rad s^{-1}
 4. 5.00 rad s^{-1}

Question ID : 41652910414

Option 1 ID : 41652941116

Option 2 ID : 41652941114

Option 3 ID : 41652941115

Option 4 ID : 41652941117

Status : Not Answered

Chosen Option : --

Q.29 A particle executes simple harmonic motion with an amplitude of 5 cm. When the particle is at 4 cm from the mean position, the magnitude of its velocity in SI units is equal to that of its acceleration. Then, its periodic time in seconds is :

- Options
1. $\frac{4\pi}{3}$
 2. $\frac{3}{8}\pi$
 3. $\frac{8\pi}{3}$

4. $\frac{7}{3} \pi$

Question ID : 41652910417

Option 1 ID : 41652941126

Option 2 ID : 41652941128

Option 3 ID : 41652941127

Option 4 ID : 41652941129

Status : Answered

Chosen Option : 3

Q.30

Two kg of a monoatomic gas is at a pressure of $4 \times 10^4 \text{ N/m}^2$. The density of the gas is 8 kg/m^3 . What is the order of energy of the gas due to its thermal motion ?

- Options
1. 10^3 J
 2. 10^5 J
 3. 10^4 J
 4. 10^6 J

Question ID : 41652910416

Option 1 ID : 41652941122

Option 2 ID : 41652941124

Option 3 ID : 41652941123

Option 4 ID : 41652941125

Status : Not Answered

Chosen Option : --

Section : Chemistry

Q.1

The ground state energy of hydrogen atom is -13.6 eV . The energy of second excited state of He^+ ion in eV is :

- Options
1. -54.4
 2. -3.4
 3. -6.04
 4. -27.2

Question ID : 41652910458

Option 1 ID : 41652941290

Option 2 ID : 41652941292

Option 3 ID : 41652941293

Option 4 ID : 41652941291

Status : Answered

Chosen Option : 2

Q.2

Haemoglobin and gold sol are examples of :

- Options
1. positively and negatively charged sols, respectively
 2. positively charged sols
 3. negatively charged sols

4. negatively and positively charged sols, respectively

Question ID : 41652910465

Option 1 ID : 41652941320

Option 2 ID : 41652941318


Option 3 ID : 41652941319

Option 4 ID : 41652941321





Status : Answered

Chosen Option : 1

- Q.3 The major product of the following reaction is :



Options



Question ID : 41652910443

Option 1 ID : 41652941231

Option 2 ID : 41652941230

Option 3 ID : 41652941233

Option 4 ID : 41652941232

Status : Answered

Chosen Option : 1

- Q.4 The amount of sugar ($C_{12}H_{22}O_{11}$) required to prepare 2 L of its 0.1 M aqueous solution is :

- Options
- 136.8 g
 - 17.1 g

3. 68.4 g

4. 34.2 g

Question ID : 41652910456

Option 1 ID : 41652941285

Option 2 ID : 41652941284

Option 3 ID : 41652941283

Option 4 ID : 41652941282

Status : Answered

Chosen Option : 3

Q.5 Among the following reactions of hydrogen with halogens, the one that requires a catalyst is :

Options 1. $\text{H}_2 + \text{I}_2 \rightarrow 2 \text{HI}$ 2. $\text{H}_2 + \text{Cl}_2 \rightarrow 2 \text{HCl}$ 3. $\text{H}_2 + \text{Br}_2 \rightarrow 2 \text{HBr}$ 4. $\text{H}_2 + \text{F}_2 \rightarrow 2 \text{HF}$

Question ID : 41652910448

Option 1 ID : 41652941253

Option 2 ID : 41652941251

Option 3 ID : 41652941252

Option 4 ID : 41652941250

Status : Answered

Chosen Option : 2

Q.6 5.1 g NH_4SH is introduced in 3.0 L evacuated flask at 327°C . 30% of the solid NH_4SH decomposed to NH_3 and H_2S as gases. The K_p of the reaction at 327°C is ($R = 0.082 \text{ L atm mol}^{-1}\text{K}^{-1}$, Molar mass of $\text{S} = 32 \text{ g mol}^{-1}$, molar mass of $\text{N} = 14 \text{ g mol}^{-1}$)

Options 1. $0.242 \times 10^{-4} \text{ atm}^2$ 2. $1 \times 10^{-4} \text{ atm}^2$ 3. $4.9 \times 10^{-3} \text{ atm}^2$ 4. 0.242 atm^2

Question ID : 41652910462

Option 1 ID : 41652941309

Option 2 ID : 41652941306

Option 3 ID : 41652941307

Option 4 ID : 41652941308

Status : Answered

Chosen Option : 1

Q.7 The reaction that is NOT involved in the ozone layer depletion mechanism in the stratosphere is :

Options 1. $\text{CF}_2\text{Cl}_2(\text{g}) \xrightarrow{\text{uv}} \dot{\text{C}}\text{Cl}(\text{g}) + \dot{\text{C}}\text{F}_2\text{Cl}(\text{g})$

2. $\text{Cl}\dot{\text{O}}(\text{g}) + \text{O}(\text{g}) \rightarrow \dot{\text{C}}\text{I}(\text{g}) + \text{O}_2(\text{g})$
3. $\text{CH}_4 + 2 \text{O}_3 \rightarrow 3 \text{CH}_2 = \text{O} + 3 \text{H}_2\text{O}$
4. $\text{HOCl}(\text{g}) \xrightarrow{h\nu} \dot{\text{O}}\text{H}(\text{g}) + \dot{\text{C}}\text{I}(\text{g})$

Question ID : 41652910455

Option 1 ID : 41652941281

Option 2 ID : 41652941279

Option 3 ID : 41652941280

Option 4 ID : 41652941278

Status : Answered

Chosen Option : 2

Q.8 In the cell

$\text{Pt}(\text{s})|\text{H}_2(\text{g}, 1\text{bar})|\text{HCl}(\text{aq})|\text{AgCl}(\text{s})|\text{Ag}(\text{s})|\text{Pt}(\text{s})$
 the cell potential is 0.92 V when a 10^{-6} molal HCl solution is used. The standard electrode potential of (AgCl/Ag,Cl⁻) electrode is :

$$\left\{ \text{Given, } \frac{2.303RT}{F} = 0.06 \text{ V at } 298 \text{ K} \right\}$$

- Options
1. 0.94 V
 2. 0.76 V
 3. 0.40 V
 4. 0.20 V

Question ID : 41652910463

Option 1 ID : 41652941313

Option 2 ID : 41652941311

Option 3 ID : 41652941310

Option 4 ID : 41652941312

Status : Answered

Chosen Option : 1

Q.9 The 71st electron of an element X with an atomic number of 71 enters into the orbital :

- Options
1. 6p
 2. 4f
 3. 5d
 4. 6s

Question ID : 41652910446

Option 1 ID : 41652941245

Option 2 ID : 41652941243

Option 3 ID : 41652941244

Option 4 ID : 41652941242

Status : Answered

Chosen Option : 3

Q.10

The correct match between item 'I' and item 'II' is :

Item 'I' (compound)	Item 'II' (reagent)
(A) Lysine	(P) 1-naphthol
(B) Furfural	(Q) ninhydrin
(C) Benzyl alcohol	(R) KMnO_4
(D) Styrene	(S) Ceric ammonium nitrate

- Options
- (A) → (Q); (B) → (P); (C) → (S); (D) → (R)
 - (A) → (Q); (B) → (P); (C) → (R); (D) → (S)
 - (A) → (R); (B) → (P); (C) → (Q); (D) → (S)
 - (A) → (Q); (B) → (R); (C) → (S); (D) → (P)

Question ID : 41652910445

Option 1 ID : 41652941240

Option 2 ID : 41652941238

Option 3 ID : 41652941239





Option 4 ID : 41652941241

Status : Answered

Chosen Option : 1

Q.11 An aromatic compound 'A' having molecular formula $\text{C}_7\text{H}_6\text{O}_2$ on treating with aqueous ammonia and heating forms compound 'B'. The compound 'B' on reaction with molecular bromine and potassium hydroxide provides compound 'C' having molecular formula $\text{C}_6\text{H}_7\text{N}$. The structure of 'A' is :

Options

- 
- 
- 
- 

Question ID : 41652910441

Option 1 ID : 41652941222

Option 2 ID : 41652941225

Option 3 ID : 41652941223

Option 4 ID : 41652941224

Status : Answered

Chosen Option : 1

Q.12 The process with negative entropy change is :

- Options
1. Dissociation of $\text{CaSO}_4(\text{s})$ to $\text{CaO}(\text{s})$ and $\text{SO}_3(\text{g})$
 2. Sublimation of dry ice
 3. Dissolution of iodine in water
 4. Synthesis of ammonia from N_2 and H_2

Question ID : 41652910460

Option 1 ID : 41652941300

Option 2 ID : 41652941298

Option 3 ID : 41652941299

Option 4 ID : 41652941301

Status : Not Answered

Chosen Option : --

Q.13 An ideal gas undergoes isothermal compression from 5 m^3 to 1 m^3 against a constant external pressure of 4 Nm^{-2} . Heat released in this process is used to increase the temperature of 1 mole of Al. If molar heat capacity of Al is $24 \text{ J mol}^{-1}\text{K}^{-1}$, the temperature of Al increases by :

- Options
1. $\frac{3}{2} \text{ K}$
 2. 2 K
 3. $\frac{2}{3} \text{ K}$
 4. 1 K

Question ID : 41652910459

Option 1 ID : 41652941297

Option 2 ID : 41652941296

Option 3 ID : 41652941295

Option 4 ID : 41652941294

Status : Answered

Chosen Option : 3

Q.14 Elevation in the boiling point for 1 molal solution of glucose is 2 K. The depression in the freezing point for 2 molal solution of glucose in the same solvent is 2 K. The relation between K_b and K_f is :

- Options
1. $K_b = 1.5 K_f$
 2. $K_b = K_f$
 3. $K_b = 0.5 K_f$
 4. $K_b = 2 K_f$

Question ID : 41652910461

Option 1 ID : 41652941302

Option 2 ID : 41652941304
 Option 3 ID : 41652941303
 Option 4 ID : 41652941305
 Status : Answered
 Chosen Option : 1

Q.15 The major product of the following reaction is :



Options

1. $\text{CH}_3\text{N}=\text{CH}-\text{CH}_2-\text{CH}_2-\text{CH}(\text{OH})-\text{CH}_2-\text{C}(=\text{O})\text{CH}_3$
2. $\text{CH}_3\text{N}=\text{CH}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}(\text{OH})-\text{C}(=\text{O})\text{CH}_3$
3. $\text{CH}_3\text{NH}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{C}(=\text{O})\text{CH}_3$
4. $\text{CH}_3\text{NH}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}=\text{CH}-\text{CH}_2-\text{CH}(\text{OH})-\text{C}(=\text{O})\text{CH}_3$


Question ID : 41652910438
 Option 1 ID : 41652941212
 Option 2 ID : 41652941211
 Option 3 ID : 41652941213
 Option 4 ID : 41652941210
 Status : Answered
 Chosen Option : 4

Q.16 Sodium metal on dissolution in liquid ammonia gives a deep blue solution due to the formation of :

- Options
1. sodium-ammonia complex
 2. sodamide
 3. sodium ion-ammonia complex
 4. ammoniated electrons

Question ID : 41652910449
 Option 1 ID : 41652941255
 Option 2 ID : 41652941254
 Option 3 ID : 41652941256
 Option 4 ID : 41652941257
 Status : Answered
 Chosen Option : 3

Q.17 For an elementary chemical reaction,



is :

- Options
1. $k_1[A_2] - k_{-1}[A]^2$
 2. $2k_1[A_2] - k_{-1}[A]^2$

3. $k_1[A_2] + k_{-1}[A]^2$
 4. $2k_1[A_2] - 2k_{-1}[A]^2$

Question ID : 41652910464
 Option 1 ID : 41652941314
 Option 2 ID : 41652941316
 Option 3 ID : 41652941317
 Option 4 ID : 41652941315
 Status : Answered
 Chosen Option : 2

Q.18 Which of the following tests cannot be used for identifying amino acids ?

- Options
1. Biuret test
 2. Barfoed test
 3. Ninhydrin test
 4. Xanthoproteic test


Question ID : 41652910444
 Option 1 ID : 41652941236
 Option 2 ID : 41652941235
 Option 3 ID : 41652941237
 Option 4 ID : 41652941234
 Status : Not Answered
 Chosen Option : --

Q.19 The difference in the number of unpaired electrons of a metal ion in its high-spin and low-spin octahedral complexes is two. The metal ion is :




- Options
1. Ni^{2+}
 2. Fe^{2+}
 3. Co^{2+}
 4. Mn^{2+}

Question ID : 41652910453
 Option 1 ID : 41652941270
 Option 2 ID : 41652941273
 Option 3 ID : 41652941272
 Option 4 ID : 41652941271
 Status : Not Answered
 Chosen Option : --

Q.20 The major product obtained in the following reaction is :



- Options
- 1.

2. 
3. 
4. 

Question ID : 41652910439

Option 1 ID : 41652941215

Option 2 ID : 41652941216

Option 3 ID : 41652941217

Option 4 ID : 41652941214

Status : Answered

Chosen Option : 3

Q.21 The pair that contains two P – H bonds in each of the oxoacids is :

- Options
1. $\text{H}_4\text{P}_2\text{O}_5$ and $\text{H}_4\text{P}_2\text{O}_6$
 2. H_3PO_2 and $\text{H}_4\text{P}_2\text{O}_5$
 3. H_3PO_3 and H_3PO_2
 4. $\text{H}_4\text{P}_2\text{O}_5$ and H_3PO_3

Question ID : 41652910450

Option 1 ID : 41652941261

Option 2 ID : 41652941260


Option 3 ID : 41652941258

Option 4 ID : 41652941259

Status : Answered

Chosen Option : 4

Q.22 Which is the most suitable reagent for the following transformation ?



- Options
1. Tollen's reagent
 2. I_2/NaOH
 3. $\text{CrO}_2\text{Cl}_2/\text{CS}_2$
 4. alkaline KMnO_4

Question ID : 41652910442

Option 1 ID : 41652941226


Option 2 ID : 41652941227

Option 3 ID : 41652941229

Option 4 ID : 41652941228

Status : **Answered**
Chosen Option : **2**

Q.23 What is the IUPAC name of the following compound ?



- Options
1. 3-Bromo-1, 2-dimethylbut-1-ene
 2. 3-Bromo-3-methyl-1, 2-dimethylprop-1-ene
 3. 2-Bromo-3-methylpent-3-ene
 4. 4-Bromo-3-methylpent-2-ene

Question ID : **41652910436**
Option 1 ID : **41652941203**
Option 2 ID : **41652941202**
Option 3 ID : **41652941205**
Option 4 ID : **41652941204**
Status : **Answered**
Chosen Option : **3**

Q.24 The number of 2-centre-2-electron and 3-centre-2-electron bonds in B_2H_6 , respectively, are :

- Options
1. 2 and 1
 2. 4 and 2
 3. 2 and 2
 4. 2 and 4

Question ID : **41652910451**
Option 1 ID : **41652941265**
Option 2 ID : **41652941263**
Option 3 ID : **41652941262**
Option 4 ID : **41652941264**
Status : **Answered**
Chosen Option : **3**

Q.25 In the reaction of oxalate with permanganate in acidic medium, the number of electrons involved in producing one molecule of CO_2 is :

- Options
1. 1
 2. 10
 3. 2
 4. 5

Question ID : **41652910452**
Option 1 ID : **41652941267**
Option 2 ID : **41652941269**

Option 3 ID : 41652941268
 Option 4 ID : 41652941266
 Status : Not Answered
 Chosen Option : --

Q.26 A reaction of cobalt(III) chloride and ethylenediamine in a 1 : 2 mole ratio generates two isomeric products A (violet coloured) and B (green coloured). A can show optical activity, but, B is optically inactive. What type of isomers does A and B represent ?

- Options
1. Geometrical isomers
 2. Coordination isomers
 3. Linkage isomers
 4. Ionisation isomers

Question ID : 41652910454
 Option 1 ID : 41652941275
 Option 2 ID : 41652941274
 Option 3 ID : 41652941277
 Option 4 ID : 41652941276
 Status : Not Answered
 Chosen Option : --

Q.27 The electrolytes usually used in the electroplating of gold and silver, respectively, are :

- Options
1. $[\text{Au}(\text{CN})_2]^-$ and $[\text{Ag}(\text{CN})_2]^-$
 2. $[\text{Au}(\text{CN})_2]^-$ and $[\text{AgCl}_2]^-$
 3. $[\text{Au}(\text{OH})_4]^-$ and $[\text{Ag}(\text{OH})_2]^-$
 4. $[\text{Au}(\text{NH}_3)_2]^+$ and $[\text{Ag}(\text{CN})_2]^-$

Question ID : 41652910447
 Option 1 ID : 41652941246
 Option 2 ID : 41652941248
 Option 3 ID : 41652941249
 Option 4 ID : 41652941247
 Status : Answered
 Chosen Option : 2

Q.28 A compound of formula A_2B_3 has the hcp lattice. Which atom forms the hcp lattice and what fraction of tetrahedral voids is occupied by the other atoms :

- Options
1. hcp lattice - A, $\frac{2}{3}$ Tetrahedral voids - B
 2. hcp lattice - A, $\frac{1}{3}$ Tetrahedral voids - B

3. hcp lattice - B, $\frac{2}{3}$ Tetrahedral voids - A
4. hcp lattice - B, $\frac{1}{3}$ Tetrahedral voids - A

Question ID : 41652910457

Option 1 ID : 41652941286

Option 2 ID : 41652941288


Option 3 ID : 41652941287

Option 4 ID : 41652941289

Status : Answered

Chosen Option : 1

Q.29 The major product of the following reaction is :



Options

- 1.
- 2.
- 3.
- 4.

Question ID : 41652910440

Option 1 ID : 41652941218

Option 2 ID : 41652941221

Option 3 ID : 41652941219


Option 4 ID : 41652941220

Status : Answered





Chosen Option : 1

Q.30

What will be the major product in the following mononitration reaction ?



Options



Question ID : 41652910437

Option 1 ID : 41652941208

Option 2 ID : 41652941207

Option 3 ID : 41652941206

Option 4 ID : 41652941209

Status : Answered

Chosen Option : 2

Section : Mathematics

Q.1 The value of λ such that sum of the squares of the roots of the quadratic equation, $x^2 + (3 - \lambda)x + 2 = \lambda$ has the least value is :

- Options
1. $\frac{15}{8}$
 2. 1
 3. $\frac{4}{9}$
 4. 2

Question ID : 41652910467

Option 1 ID : 41652941329
 Option 2 ID : 41652941328
 Option 3 ID : 41652941326
 Option 4 ID : 41652941327
 Status : Answered
 Chosen Option : 3

Q.2 The value of

$$\cos \frac{\pi}{2^2} \cdot \cos \frac{\pi}{2^3} \cdot \dots \cdot \cos \frac{\pi}{2^{10}} \cdot \sin \frac{\pi}{2^{10}}$$

is :

Options

1. $\frac{1}{512}$

2. $\frac{1}{1024}$

3. $\frac{1}{256}$

4. $\frac{1}{2}$

Question ID : 41652910493
 Option 1 ID : 41652941432
 Option 2 ID : 41652941430
 Option 3 ID : 41652941433
 Option 4 ID : 41652941431
 Status : Answered
 Chosen Option : 1

Q.3 The curve amongst the family of curves represented by the differential equation, $(x^2 - y^2)dx + 2xy dy = 0$ which passes through (1, 1), is :

- Options
1. a circle with centre on the x -axis.
 2. an ellipse with major axis along the y -axis.
 3. a circle with centre on the y -axis.
 4. a hyperbola with transverse axis along the x -axis.

Question ID : 41652910482
 Option 1 ID : 41652941386
 Option 2 ID : 41652941388
 Option 3 ID : 41652941387
 Option 4 ID : 41652941389
 Status : Not Answered
 Chosen Option : --

Q.4 Let $f: (-1, 1) \rightarrow \mathbb{R}$ be a function defined by

$$f(x) = \max \left\{ -|x|, -\sqrt{1-x^2} \right\}. \text{ If } K \text{ be}$$

the set of all points at which f is not differentiable, then K has exactly :

- Options
1. five elements
 2. one element
 3. three elements
 4. two elements

Question ID : 41652910476
 Option 1 ID : 41652941365
 Option 2 ID : 41652941362
 Option 3 ID : 41652941364
 Option 4 ID : 41652941363
 Status : Answered
 Chosen Option : 3

Q.5 The positive value of λ for which the co-efficient of x^2 in the expression

$$x^2 \left(\sqrt{x} + \frac{\lambda}{x^2} \right)^{10} \text{ is } 720, \text{ is :}$$

- Options
1. 4
 2. $2\sqrt{2}$
 3. $\sqrt{5}$
 4. 3

Question ID : 41652910472
 Option 1 ID : 41652941349
 Option 2 ID : 41652941347
 Option 3 ID : 41652941346
 Option 4 ID : 41652941348
 Status : Not Answered
 Chosen Option : --

Q.6 The tangent to the curve, $y = xe^{x^2}$ passing through the point $(1, e)$ also passes through the point :

- Options
1. $(2, 3e)$
 2. $\left(\frac{4}{3}, 2e \right)$
 3. $\left(\frac{5}{3}, 2e \right)$
 4. $(3, 6e)$

Question ID : 41652910477
 Option 1 ID : 41652941366
 Option 2 ID : 41652941368
 Option 3 ID : 41652941369
 Option 4 ID : 41652941367
 Status : Answered
 Chosen Option : 2

Q.7

Let \mathbb{N} be the set of natural numbers and two functions f and g be defined as $f, g : \mathbb{N} \rightarrow \mathbb{N}$ such that

$$f(n) = \begin{cases} \frac{n+1}{2} & \text{if } n \text{ is odd} \\ \frac{n}{2} & \text{if } n \text{ is even} \end{cases}$$

and $g(n) = n - (-1)^n$. Then $f \circ g$ is :

- Options
1. onto but not one-one.
 2. one-one but not onto.
 3. both one-one and onto.
 4. neither one-one nor onto.

Question ID : 41652910466

Option 1 ID : 41652941324

Option 2 ID : 41652941323

Option 3 ID : 41652941322

Option 4 ID : 41652941325

Status : Answered

Chosen Option : 2

Q.8 The number of values of $\theta \in (0, \pi)$ for which the system of linear equations

$$x + 3y + 7z = 0$$

$$-x + 4y + 7z = 0$$

$$(\sin 3\theta)x + (\cos 2\theta)y + 2z = 0$$

has a non-trivial solution, is :

- Options
1. three
 2. two
 3. four
 4. one

Question ID : 41652910470

Option 1 ID : 41652941339

Option 2 ID : 41652941340

Option 3 ID : 41652941338

Option 4 ID : 41652941341

Status : Not Answered

Chosen Option : --

Q.9

Let $\vec{\alpha} = (\lambda - 2)\vec{a} + \vec{b}$ and

$\vec{\beta} = (4\lambda - 2)\vec{a} + 3\vec{b}$ be two given

vectors where vectors \vec{a} and \vec{b} are non-collinear. The value of λ for which

vectors $\vec{\alpha}$ and $\vec{\beta}$ are collinear, is :

- Options
1. -4

2. -3
3. 4
4. 3

Question ID : 41652910490
 Option 1 ID : 41652941421
 Option 2 ID : 41652941418
 Option 3 ID : 41652941420
 Option 4 ID : 41652941419
 Status : Answered
 Chosen Option : 2

Q.10 Two sides of a parallelogram are along the lines, $x + y = 3$ and $x - y + 3 = 0$. If its diagonals intersect at $(2, 4)$, then one of its vertex is :

- Options
1. $(3, 5)$
 2. $(2, 1)$
 3. $(2, 6)$
 4. $(3, 6)$

Question ID : 41652910484
 Option 1 ID : 41652941397
 Option 2 ID : 41652941395
 Option 3 ID : 41652941394
 Option 4 ID : 41652941396
 Status : Answered
 Chosen Option : 3

Q.11

If $\int_0^x f(t) dt = x^2 + \int_x^1 t^2 f(t) dt$, then

$f'(1/2)$ is :

- Options
1. $\frac{24}{25}$
 2. $\frac{18}{25}$
 3. $\frac{4}{5}$
 4. $\frac{6}{25}$

Question ID : 41652910481
 Option 1 ID : 41652941384
 Option 2 ID : 41652941382
 Option 3 ID : 41652941383
 Option 4 ID : 41652941385
 Status : Answered
 Chosen Option : 2

Q.12

Let $z = \left(\frac{\sqrt{3}}{2} + \frac{i}{2}\right)^5 + \left(\frac{\sqrt{3}}{2} - \frac{i}{2}\right)^5$. If $R(z)$

and $I(z)$ respectively denote the real and imaginary parts of z , then :

- Options
1. $I(z) = 0$
 2. $R(z) > 0$ and $I(z) > 0$
 3. $R(z) < 0$ and $I(z) > 0$
 4. $R(z) = -3$

Question ID : 41652910468

Option 1 ID : 41652941333

Option 2 ID : 41652941330

Option 3 ID : 41652941331

Option 4 ID : 41652941332

Status : Answered

Chosen Option : 2

Q.13 If the probability of hitting a target by a shooter, in any shot, is $\frac{1}{3}$, then the minimum number of independent shots at the target required by him so that the probability of hitting the target at least once is greater than $\frac{5}{6}$, is :

- Options
1. 3
 2. 6
 3. 5
 4. 4

Question ID : 41652910492

Option 1 ID : 41652941426

Option 2 ID : 41652941429

Option 3 ID : 41652941428

Option 4 ID : 41652941427

Status : Answered

Chosen Option : 1

Q.14 If $\int x^5 e^{-4x^3} dx = \frac{1}{48} e^{-4x^3} f(x) + C$, where C is a constant of integration, then $f(x)$ is equal to :

- Options
1. $-2x^3 - 1$
 2. $-4x^3 - 1$
 3. $-2x^3 + 1$
 4. $4x^3 + 1$

Question ID : 41652910479

Option 1 ID : 41652941374

Option 2 ID : 41652941375

Option 3 ID : 41652941377

Option 4 ID : 41652941376
 Status : Answered
 Chosen Option : 4

Q.15 If the area of an equilateral triangle inscribed in the circle, $x^2 + y^2 + 10x + 12y + c = 0$ is $27\sqrt{3}$ sq. units then c is equal to :

- Options
1. 13
 2. 20
 3. -25
 4. 25

Question ID : 41652910485
 Option 1 ID : 41652941398
 Option 2 ID : 41652941399
 Option 3 ID : 41652941401
 Option 4 ID : 41652941400
 Status : Answered
 Chosen Option : 3

Q.16 Consider the following three statements :

P : 5 is a prime number.
 Q : 7 is a factor of 192.
 R : L.C.M. of 5 and 7 is 35.
 Then the truth value of which one of the following statements is true ?

- Options
1. $(\sim P) \vee (Q \wedge R)$
 2. $(P \wedge Q) \vee (\sim R)$
 3. $(\sim P) \wedge (\sim Q \wedge R)$
 4. $P \vee (\sim Q \wedge R)$

Question ID : 41652910495
 Option 1 ID : 41652941441
 Option 2 ID : 41652941438
 Option 3 ID : 41652941439
 Option 4 ID : 41652941440
 Status : Answered
 Chosen Option : 1

Q.17 The length of the chord of the parabola $x^2 = 4y$ having equation $x - \sqrt{2}y + 4\sqrt{2} = 0$ is :

- Options
1. $3\sqrt{2}$
 2. $2\sqrt{11}$
 3. $8\sqrt{2}$
 4. $6\sqrt{3}$

Question ID : 41652910487
 Option 1 ID : 41652941406

Option 2 ID : 41652941407
 Option 3 ID : 41652941409
 Option 4 ID : 41652941408
 Status : Not Answered
 Chosen Option : --

Q.18

Let $A = \begin{bmatrix} 2 & b & 1 \\ b & b^2 + 1 & b \\ 1 & b & 2 \end{bmatrix}$ where $b > 0$. Then

the minimum value of $\frac{\det(A)}{b}$ is :

- Options
1. $2\sqrt{3}$
 2. $-2\sqrt{3}$
 3. $-\sqrt{3}$
 4. $\sqrt{3}$

Question ID : 41652910469
 Option 1 ID : 41652941337
 Option 2 ID : 41652941334
 Option 3 ID : 41652941335
 Option 4 ID : 41652941336
 Status : Not Answered
 Chosen Option : --

Q.19 Let

$$S = \left\{ (x, y) \in \mathbb{R}^2 : \frac{y^2}{1+r} - \frac{x^2}{1-r} = 1 \right\},$$

where $r \neq \pm 1$. Then S represents :

Options a hyperbola whose eccentricity is

1. $\frac{2}{\sqrt{1-r}}$, when $0 < r < 1$.

an ellipse whose eccentricity is

2. $\sqrt{\frac{2}{r+1}}$, when $r > 1$.

a hyperbola whose eccentricity is

3. $\frac{2}{\sqrt{r+1}}$, when $0 < r < 1$.

an ellipse whose eccentricity is

4. $\frac{1}{\sqrt{r+1}}$, when $r > 1$.

Question ID : 41652910486
 Option 1 ID : 41652941405
 Option 2 ID : 41652941402
 Option 3 ID : 41652941404
 Option 4 ID : 41652941403
 Status : Answered

Chosen Option : 3

Q.20

$$\text{If } \sum_{r=0}^{25} \left\{ {}^{50}C_r \cdot 50^{-r} C_{25-r} \right\} = K \left({}^{50}C_{25} \right),$$

then K is equal to :

- Options
1. $(25)^2$
 2. $2^{25} - 1$
 3. 2^{24}
 4. 2^{25}

Question ID : 41652910471

Option 1 ID : 41652941342

Option 2 ID : 41652941344

Option 3 ID : 41652941343

Option 4 ID : 41652941345

Status : Answered

Chosen Option : 3

Q.21

The plane which bisects the line segment joining the points $(-3, -3, 4)$ and $(3, 7, 6)$ at right angles, passes through which one of the following points ?

- Options
1. $(-2, 3, 5)$
 2. $(4, -1, 7)$
 3. $(2, 1, 3)$
 4. $(4, 1, -2)$

Question ID : 41652910488

Option 1 ID : 41652941413

Option 2 ID : 41652941410

Option 3 ID : 41652941412

Option 4 ID : 41652941411

Status : Answered

Chosen Option : 4

Q.22

$$\text{The value of } \cot \left(\sum_{n=1}^{19} \cot^{-1} \left(1 + \sum_{p=1}^n 2p \right) \right)$$

is :

- Options
1. $\frac{21}{19}$
 2. $\frac{19}{21}$
 3. $\frac{22}{23}$
 4. $\frac{23}{22}$

Question ID : 41652910474

Option 1 ID : 41652941356

Option 2 ID : 41652941357
 Option 3 ID : 41652941354
 Option 4 ID : 41652941355
 Status : Answered
 Chosen Option : 2

Q.23 If mean and standard deviation of 5 observations x_1, x_2, x_3, x_4, x_5 are 10 and 3, respectively, then the variance of 6 observations x_1, x_2, \dots, x_5 and -50 is equal to :

- Options
1. 509.5
 2. 586.5
 3. 582.5
 4. 507.5

Question ID : 41652910491
 Option 1 ID : 41652941424
 Option 2 ID : 41652941425
 Option 3 ID : 41652941423
 Option 4 ID : 41652941422
 Status : Not Answered
 Chosen Option : --

Q.24 Let f be a differentiable function such that

$$f'(x) = 7 - \frac{3}{4} \frac{f(x)}{x}, (x > 0) \quad \text{and}$$

$$f(1) \neq 4. \text{ Then } \lim_{x \rightarrow 0^+} x f\left(\frac{1}{x}\right):$$

- Options
1. exists and equals $\frac{4}{7}$.
 2. exists and equals 4.
 3. does not exist.
 4. exists and equals 0.

Question ID : 41652910475
 Option 1 ID : 41652941360
 Option 2 ID : 41652941359
 Option 3 ID : 41652941358
 Option 4 ID : 41652941361
 Status : Not Answered
 Chosen Option : --

Q.25 Two vertices of a triangle are (0, 2) and (4, 3). If its orthocentre is at the origin, then its third vertex lies in which quadrant ?

- Options
1. third
 2. second
 3. first
 4. fourth

Question ID : 41652910483
 Option 1 ID : 41652941392
 Option 2 ID : 41652941391
 Option 3 ID : 41652941390
 Option 4 ID : 41652941393
 Status : Not Answered
 Chosen Option : --

Q.26

The value of $\int_{-\pi/2}^{\pi/2} \frac{dx}{[x] + [\sin x] + 4}$, where

$[t]$ denotes the greatest integer less than or equal to t , is :

Options

1. $\frac{1}{12} (7\pi + 5)$
2. $\frac{1}{12} (7\pi - 5)$
3. $\frac{3}{20} (4\pi - 3)$
4. $\frac{3}{10} (4\pi - 3)$

Question ID : 41652910480
 Option 1 ID : 41652941380
 Option 2 ID : 41652941381
 Option 3 ID : 41652941378
 Option 4 ID : 41652941379
 Status : Not Answered
 Chosen Option : --

Q.27 On which of the following lines lies the point of intersection of the line,

$\frac{x-4}{2} = \frac{y-5}{2} = \frac{z-3}{1}$ and the plane,
 $x+y+z=2$?

Options

1. $\frac{x+3}{3} = \frac{4-y}{3} = \frac{z+1}{-2}$
2. $\frac{x-4}{1} = \frac{y-5}{1} = \frac{z-5}{-1}$
3. $\frac{x-1}{1} = \frac{y-3}{2} = \frac{z+4}{-5}$
4. $\frac{x-2}{2} = \frac{y-3}{2} = \frac{z+3}{3}$

Question ID : 41652910489
 Option 1 ID : 41652941417
 Option 2 ID : 41652941415
 Option 3 ID : 41652941414
 Option 4 ID : 41652941416

Status : **Answered**
Chosen Option : **3**

Q.28 Let $a_1, a_2, a_3, \dots, a_{10}$ be in G.P. with $a_i > 0$ for $i = 1, 2, \dots, 10$ and S be the set of pairs (r, k) , $r, k \in \mathbb{N}$ (the set of natural numbers) for which

$$\begin{vmatrix} \log_e a_1^r a_2^k & \log_e a_2^r a_3^k & \log_e a_3^r a_4^k \\ \log_e a_4^r a_5^k & \log_e a_5^r a_6^k & \log_e a_6^r a_7^k \\ \log_e a_7^r a_8^k & \log_e a_8^r a_9^k & \log_e a_9^r a_{10}^k \end{vmatrix} = 0$$

Then the number of elements in S , is :

Options 1. 4

2. infinitely many

3. 2

4. 10

Question ID : **41652910473**
Option 1 ID : **41652941351**
Option 2 ID : **41652941353**
Option 3 ID : **41652941350**
Option 4 ID : **41652941352**
Status : **Not Answered**
Chosen Option : --

Q.29 With the usual notation, in ΔABC , if $\angle A + \angle B = 120^\circ$, $a = \sqrt{3} + 1$ and $b = \sqrt{3} - 1$, then the ratio $\angle A : \angle B$, is :

Options 1. 7 : 1

2. 5 : 3

3. 9 : 7

4. 3 : 1

Question ID : **41652910494**
Option 1 ID : **41652941437**
Option 2 ID : **41652941434**
Option 3 ID : **41652941436**
Option 4 ID : **41652941435**
Status : **Answered**
Chosen Option : **2**

Q.30 A helicopter is flying along the curve given by $y - x^{3/2} = 7$, ($x \geq 0$). A soldier positioned at the point $\left(\frac{1}{2}, 7\right)$ wants to shoot down the helicopter when it is nearest to him. Then this nearest distance is :

Options 1. $\frac{\sqrt{5}}{6}$

2. $\frac{1}{3}\sqrt{\frac{7}{3}}$

3. $\frac{1}{6}\sqrt{\frac{7}{3}}$

4. $\frac{1}{2}$

Question ID : 41652910478

Option 1 ID : 41652941370

Option 2 ID : 41652941372

Option 3 ID : 41652941373

Option 4 ID : 41652941371

Status : Answered

Chosen Option : 2