





JEE MAIN 2019

Application No	
Candidate Name	
Roll No.	
Test Date	12/01/2019
Test Time	2:30 PM - 5:30 PM
Subject	Paper I EH

Section : Physics

Q.1 The moment of inertia of a solid sphere, about an axis parallel to its diameter and at a distance of x from it, is ' $I(x)$ '. Which one of the graphs represents the variation of $I(x)$ with x correctly ?

Options



Question ID : **4165299871**
 Option 1 ID : **41652938944**
 Option 2 ID : **41652938943**
 Option 3 ID : **41652938942**
 Option 4 ID : **41652938945**

Status : **Answered**

Chosen Option : **2**


Q.2 A load of mass M kg is suspended from a steel wire of length 2 m and radius 1.0 mm in Searle's apparatus experiment. The increase in length produced in the wire is 4.0 mm. Now the load is fully immersed in a liquid of relative density 2. The relative density of the material of load is 8.

The new value of increase in length of the steel wire is :

- Options
1. 3.0 mm
 2. 4.0 mm
 3. 5.0 mm
 4. zero

Question ID : 4165299894
 Option 1 ID : 41652939035
 Option 2 ID : 41652939034
 Option 3 ID : 41652939037
 Option 4 ID : 41652939036
 Status : Answered
 Chosen Option : 4

Q.3



In the above circuit, $C = \frac{\sqrt{3}}{2} \mu\text{F}$, $R_2 = 20 \Omega$,

$L = \frac{\sqrt{3}}{10} \text{ H}$ and $R_1 = 10 \Omega$. Current in

L- R_1 path is I_1 and in C- R_2 path it is I_2 . The voltage of A.C source is given by,

$V = 200\sqrt{2}\sin(100t)$ volts. The phase difference between I_1 and I_2 is :

- Options
1. 60°
 2. 30°
 3. 90°
 4. 0°

Question ID : 4165299882
 Option 1 ID : 41652938987
 Option 2 ID : 41652938989
 Option 3 ID : 41652938988
 Option 4 ID : 41652938986
 Status : Not Answered
 Chosen Option : --

Q.4


An ideal gas is enclosed in a cylinder at pressure of 2 atm and temperature, 300 K. The mean time between two successive collisions is $6 \times 10^{-8} \text{ s}$. If the pressure is doubled and temperature is increased to 500 K, the mean time between two successive collisions will be close to :

- Options
1. $2 \times 10^{-7} \text{ s}$

- 2. $4 \times 10^{-8} \text{ s}$
- 3. $0.5 \times 10^{-8} \text{ s}$
- 4. $3 \times 10^{-6} \text{ s}$

Question ID : 4165299876
 Option 1 ID : 41652938963
 Option 2 ID : 41652938965
 Option 3 ID : 41652938964
 Option 4 ID : 41652938962
 Status : Not Answered
 Chosen Option : --

Q.5




In the figure, given that V_{BB} supply can vary from 0 to 5.0 V, $V_{CC} = 5 \text{ V}$, $\beta_{dc} = 200$, $R_B = 100 \text{ k}\Omega$, $R_C = 1 \text{ k}\Omega$ and $V_{BE} = 1.0 \text{ V}$, The minimum base current and the input voltage at which the transistor will go to saturation, will be, respectively :

- Options
- 1. $25 \mu\text{A}$ and 3.5 V
 - 2. $20 \mu\text{A}$ and 3.5 V
 - 3. $25 \mu\text{A}$ and 2.8 V
 - 4. $20 \mu\text{A}$ and 2.8 V

Question ID : 4165299892
 Option 1 ID : 41652939026
 Option 2 ID : 41652939029
 Option 3 ID : 41652939028
 Option 4 ID : 41652939027
 Status : Not Answered
 Chosen Option : --

Q.6 In the circuit shown, find C if the effective capacitance of the whole circuit is to be $0.5 \mu\text{F}$. All values in the circuit are in μF .



- Options
- 1. $\frac{7}{11} \mu\text{F}$

2. $\frac{6}{5} \mu\text{F}$

3. $4 \mu\text{F}$

4. $\frac{7}{10} \mu\text{F}$

Question ID : 4165299880

Option 1 ID : 41652938980

Option 2 ID : 41652938981

Option 3 ID : 41652938979

Option 4 ID : 41652938978

Status : Not Answered

Chosen Option : --

Q.7 An alpha-particle of mass m suffers 1-dimensional elastic collision with a nucleus at rest of unknown mass. It is scattered directly backwards losing, 64% of its initial kinetic energy. The mass of the nucleus is :

- Options
1. $2m$
 2. $3.5m$
 3. $1.5m$
 4. $4m$

Question ID : 4165299869

Option 1 ID : 41652938936

Option 2 ID : 41652938935

Option 3 ID : 41652938937

Option 4 ID : 41652938934

Status : Not Answered

Chosen Option : --

Q.8 A 10 m long horizontal wire extends from North East to South West. It is falling with a speed of 5.0 ms^{-1} , at right angles to the horizontal component of the earth's magnetic field, of $0.3 \times 10^{-4} \text{ Wb/m}^2$. The value of the induced emf in wire is :

- Options
1. $1.5 \times 10^{-3} \text{ V}$
 2. $1.1 \times 10^{-3} \text{ V}$
 3. $2.5 \times 10^{-3} \text{ V}$
 4. $0.3 \times 10^{-3} \text{ V}$

Question ID : 4165299885

Option 1 ID : 41652939000

Option 2 ID : 41652938998

Option 3 ID : 41652938999

Option 4 ID : 41652939001

Status : Answered

Chosen Option : 4

Q.9

To double the covering range of a TV transmission tower, its height should be multiplied by :

- Options
1. $\frac{1}{\sqrt{2}}$
 2. 2
 3. 4
 4. $\sqrt{2}$

Question ID : 4165299893
 Option 1 ID : 41652939033
 Option 2 ID : 41652939030
 Option 3 ID : 41652939031
 Option 4 ID : 41652939032
 Status : Answered
 Chosen Option : 4

Q.10 A plano-convex lens (focal length f_2 , refractive index μ_2 , radius of curvature R) fits exactly into a plano-concave lens (focal length f_1 , refractive index μ_1 , radius of curvature R). Their plane surfaces are parallel to each other. Then, the focal length of the combination will be :

- Options
1. $f_1 - f_2$
 2. $\frac{R}{\mu_2 - \mu_1}$
 3. $\frac{2f_1 f_2}{f_1 + f_2}$
 4. $f_1 + f_2$

Question ID : 4165299887
 Option 1 ID : 41652939007
 Option 2 ID : 41652939009
 Option 3 ID : 41652939008
 Option 4 ID : 41652939006
 Status : Answered
 Chosen Option : 4

Q.11 A vertical closed cylinder is separated into two parts by a frictionless piston of mass m and of negligible thickness. The piston is free to move along the length of the cylinder. The length of the cylinder above the piston is l_1 , and that below the piston is l_2 , such that $l_1 > l_2$. Each part of the cylinder contains n moles of an ideal gas at equal temperature T . If the piston is stationary, its mass, m , will be given by :

(R is universal gas constant and g is the acceleration due to gravity)

Options

1. $\frac{RT}{ng} \left[\frac{l_1 - 3l_2}{l_1 l_2} \right]$

2. $\frac{RT}{g} \left[\frac{2l_1 + l_2}{l_1 l_2} \right]$

3. $\frac{nRT}{g} \left[\frac{1}{l_2} + \frac{1}{l_1} \right]$

4. $\frac{nRT}{g} \left[\frac{l_1 - l_2}{l_1 l_2} \right]$

Question ID : 4165299875

Option 1 ID : 41652938961

Option 2 ID : 41652938960

Option 3 ID : 41652938958

Option 4 ID : 41652938959

Status : Answered

Chosen Option : 4

Q.12 Two satellites, A and B, have masses m and $2m$ respectively. A is in a circular orbit of radius R , and B is in a circular orbit of radius $2R$ around the earth. The ratio of their kinetic energies, T_A/T_B , is :

Options

1. $\frac{1}{2}$

2. 1

3. 2

4. $\sqrt{\frac{1}{2}}$

Question ID : 4165299872

Option 1 ID : 41652938947

Option 2 ID : 41652938948

Option 3 ID : 41652938946

Option 4 ID : 41652938949

Status : Answered

Chosen Option : 1

Q.13 A long cylindrical vessel is half filled with a liquid. When the vessel is rotated about its own vertical axis, the liquid rises up near the wall. If the radius of vessel is 5 cm and its rotational speed is 2 rotations per second, then the difference in the heights between the centre and the sides, in cm, will be :

Options

1. 2.0


2. 0.1

3. 0.4

4. 1.2

Question ID : 4165299873
 Option 1 ID : 41652938952
 Option 2 ID : 41652938951
 Option 3 ID : 41652938950
 Option 4 ID : 41652938953
 Status : Answered
 Chosen Option : 1

- Q.14 A block kept on a rough inclined plane, as shown in the figure, remains at rest upto a maximum force 2 N down the inclined plane. The maximum external force up the inclined plane that does not move the block is 10 N. The coefficient of static friction between the block and the plane is :
 [Take $g = 10 \text{ m/s}^2$]



Options

1. $\frac{\sqrt{3}}{2}$
2. $\frac{\sqrt{3}}{4}$
3. $\frac{1}{2}$
4. $\frac{2}{3}$

Question ID : 4165299868
 Option 1 ID : 41652938933
 Option 2 ID : 41652938931
 Option 3 ID : 41652938930
 Option 4 ID : 41652938932
 Status : Answered
 Chosen Option : 1


- Q.15 In a Frank-Hertz experiment, an electron of energy 5.6 eV passes through mercury vapour and emerges with an energy 0.7 eV. The minimum wavelength of photons emitted by mercury atoms is close to :

- Options
1. 1700 nm
 2. 2020 nm
 3. 220 nm
 4. 250 nm

Question ID : 4165299890
 Option 1 ID : 41652939020
 Option 2 ID : 41652939019
 Option 3 ID : 41652939018
 Option 4 ID : 41652939021
 Status : Answered

Chosen Option : 2

- Q.16** A particle of mass 20 g is released with an initial velocity 5 m/s along the curve from the point A, as shown in the figure. The point A is at height h from point B. The particle slides along the frictionless surface. When the particle reaches point B, its angular momentum about O will be :
(Take $g = 10 \text{ m/s}^2$)



- Options**
1. $2 \text{ kg-m}^2/\text{s}$
 2. $8 \text{ kg-m}^2/\text{s}$
 3. $6 \text{ kg-m}^2/\text{s}$
 4. $3 \text{ kg-m}^2/\text{s}$

Question ID : 4165299870

Option 1 ID : 41652938940

Option 2 ID : 41652938939

Option 3 ID : 41652938938

Option 4 ID : 41652938941

Status : Not Answered

Chosen Option : --

- Q.17** A galvanometer, whose resistance is 50 ohm, has 25 divisions in it. When a current of $4 \times 10^{-4} \text{ A}$ passes through it, its needle (pointer) deflects by one division. To use this galvanometer as a voltmeter of range 2.5 V, it should be connected to a resistance of :

- Options**
1. 250 ohm
 2. 200 ohm
 3. 6200 ohm
 4. 6250 ohm

Question ID : 4165299883

Option 1 ID : 41652938992

Option 2 ID : 41652938993

Option 3 ID : 41652938991

Option 4 ID : 41652938990





Status : Not Answered

Chosen Option : --

Q.18

A soap bubble, blown by a mechanical pump at the mouth of a tube, increases in volume, with time, at a constant rate. The graph that correctly depicts the time dependence of pressure inside the bubble is given by :

Options



Question ID : 4165299874

Option 1 ID : 41652938956

Option 2 ID : 41652938957


Option 3 ID : 41652938955

Option 4 ID : 41652938954

Status : Answered

Chosen Option : 3

Q.19 In the given circuit diagram, the currents, $I_1 = -0.3$ A, $I_4 = 0.8$ A and $I_5 = 0.4$ A, are flowing as shown. The currents I_2 , I_3 and I_6 , respectively, are :



- Options
1. 1.1 A , -0.4 A, 0.4 A
 2. 1.1 A , 0.4 A, 0.4 A

3. 0.4 A , 1.1 A, 0.4 A
4. -0.4 A , 0.4 A, 1.1 A

Question ID : 4165299881
 Option 1 ID : 41652938982
 Option 2 ID : 41652938984
 Option 3 ID : 41652938985
 Option 4 ID : 41652938983

Status : Answered

Chosen Option : 2

Q.20

A resonance tube is old and has jagged end. It is still used in the laboratory to determine velocity of sound in air. A tuning fork of frequency 512 Hz produces first resonance when the tube is filled with water to a mark 11 cm below a reference mark, near the open end of the tube. The experiment is repeated with another fork of frequency 256 Hz which produces first resonance when water reaches a mark 27 cm below the reference mark. The velocity of sound in air, obtained in the experiment, is close to :

- Options
1. 322 ms^{-1}
 2. 341 ms^{-1}
 3. 335 ms^{-1}
 4. 328 ms^{-1}

Question ID : 4165299878
 Option 1 ID : 41652938972
 Option 2 ID : 41652938973
 Option 3 ID : 41652938971
 Option 4 ID : 41652938970

Status : Not Answered

Chosen Option : --

Q.21

A paramagnetic material has 10^{28} atoms/ m^3 . Its magnetic susceptibility at temperature 350 K is 2.8×10^{-4} . Its susceptibility at 300 K is :

- Options
1. 3.267×10^{-4}
 2. 3.672×10^{-4}
 3. 3.726×10^{-4}
 4. 2.672×10^{-4}

Question ID : 4165299884
 Option 1 ID : 41652938995
 Option 2 ID : 41652938997
 Option 3 ID : 41652938994
 Option 4 ID : 41652938996

Status : Answered

Chosen Option : 2

Q.22

In a radioactive decay chain, the initial nucleus is ${}_{90}^{232}\text{Th}$. At the end there are 6 α -particles and 4 β -particles which are emitted. If the end nucleus is ${}^A_Z\text{X}$, A and Z are given by :

- Options
1. $A = 208 ; Z = 80$
 2. $A = 202 ; Z = 80$
 3. $A = 208 ; Z = 82$
 4. $A = 200 ; Z = 81$

Question ID : 4165299891
 Option 1 ID : 41652939024
 Option 2 ID : 41652939022
 Option 3 ID : 41652939023
 Option 4 ID : 41652939025

Status : Answered

Chosen Option : 1

Q.23 Let l , r , c and v represent inductance, resistance, capacitance and voltage, respectively. The dimension of $\frac{l}{rcv}$ in SI units will be :


- Options
1. $[LA^{-2}]$
 2. $[A^{-1}]$
 3. $[LTA]$
 4. $[LT^2]$

Question ID : 4165299866
 Option 1 ID : 41652938925
 Option 2 ID : 41652938923
 Option 3 ID : 41652938922
 Option 4 ID : 41652938924

Status : Answered

Chosen Option : 1

Q.24 Formation of real image using a biconvex lens is shown below :



If the whole set up is immersed in water without disturbing the object and the screen positions, what will one observe on the screen ?

- Options
1. Image disappears
 2. Magnified image

3. Erect real image
4. No change

Question ID : 4165299888
 Option 1 ID : 41652939012
 Option 2 ID : 41652939011
 Option 3 ID : 41652939013
 Option 4 ID : 41652939010
 Status : Answered
 Chosen Option : 2

Q.25 When a certain photosensitive surface is illuminated with monochromatic light of frequency ν , the stopping potential for the photo current is $-V_0/2$. When the surface is illuminated by monochromatic light of frequency $\nu/2$, the stopping potential is $-V_0$. The threshold frequency for photoelectric emission is :

- Options
1. $\frac{5\nu}{3}$
 2. $\frac{4}{3}\nu$
 3. 2ν
 4. $\frac{3\nu}{2}$

Question ID : 4165299889
 Option 1 ID : 41652939015
 Option 2 ID : 41652939016
 Option 3 ID : 41652939014
 Option 4 ID : 41652939017
 Status : Answered
 Chosen Option : 4

Q.26 A simple harmonic motion is represented by :

$$y = 5(\sin 3\pi t + \sqrt{3} \cos 3\pi t) \text{ cm}$$

The amplitude and time period of the motion are :

- Options
1. 10 cm, $\frac{2}{3}$ s
 2. 10 cm, $\frac{3}{2}$ s
 3. 5 cm, $\frac{3}{2}$ s
 4. 5 cm, $\frac{2}{3}$ s

Question ID : 4165299877
 Option 1 ID : 41652938969
 Option 2 ID : 41652938968

Option 3 ID : 41652938966


Option 4 ID : 41652938967

Status : Answered

Chosen Option : 2

Q.27

Two particles A, B are moving on two concentric circles of radii R_1 and R_2 with equal angular speed ω . At $t=0$, their positions and direction of motion are shown in the figure :



The relative velocity $\vec{v}_A - \vec{v}_B$ at $t = \frac{\pi}{2\omega}$ is given by :

Options

1. $\omega(R_1 + R_2)\hat{i}$
2. $-\omega(R_1 + R_2)\hat{i}$
3. $\omega(R_2 - R_1)\hat{i}$
4. $\omega(R_1 - R_2)\hat{i}$

Question ID : 4165299867

Option 1 ID : 41652938929

Option 2 ID : 41652938928

Option 3 ID : 41652938927

Option 4 ID : 41652938926

Status : Not Answered

Chosen Option : --

Q.28

The mean intensity of radiation on the surface of the Sun is about 10^8 W/m^2 . The rms value of the corresponding magnetic field is closest to :

Options

1. 1 T
2. 10^2 T
3. 10^{-2} T
4. 10^{-4} T

Question ID : 4165299886

Option 1 ID : 41652939004

Option 2 ID : 41652939005

Option 3 ID : 41652939003

Option 4 ID : 41652939002

Status : **Not Answered**
Chosen Option : --


Q.29 A parallel plate capacitor with plates of area 1 m^2 each, are at a separation of 0.1 m . If the electric field between the plates is 100 N/C , the magnitude of charge on each plate is :

(Take $\epsilon_0 = 8.85 \times 10^{-12} \frac{\text{C}^2}{\text{N-m}^2}$)

- Options**
- $7.85 \times 10^{-10} \text{ C}$
 - $6.85 \times 10^{-10} \text{ C}$
 - $8.85 \times 10^{-10} \text{ C}$
 - $9.85 \times 10^{-10} \text{ C}$

Question ID : **4165299879**
Option 1 ID : **41652938975**
Option 2 ID : **41652938974**
Option 3 ID : **41652938976**
Option 4 ID : **41652938977**
Status : **Not Answered**
Chosen Option : --

Q.30 The charge on a capacitor plate in a circuit, as a function of time, is shown in the figure :



What is the value of current at $t = 4 \text{ s}$?

- Options**
- zero
 - $3 \mu\text{A}$
 - $2 \mu\text{A}$
 - $1.5 \mu\text{A}$

Question ID : **4165299895**
Option 1 ID : **41652939041**
Option 2 ID : **41652939038**
Option 3 ID : **41652939040**
Option 4 ID : **41652939039**
Status : **Answered**
Chosen Option : **1**

Section : Chemistry

Q.1 8 g of NaOH is dissolved in 18 g of H_2O . Mole fraction of NaOH in solution and molality (in mol kg^{-1}) of the solution respectively are :

- Options
1. 0.2, 22.20
 2. 0.2, 11.11
 3. 0.167, 11.11
 4. 0.167, 22.20

Question ID : 4165299916
Option 1 ID : 41652939125
Option 2 ID : 41652939123
Option 3 ID : 41652939122
Option 4 ID : 41652939124
Status : Answered
Chosen Option : 1

Q.2 The magnetic moment of an octahedral homoleptic Mn(II) complex is 5.9 BM. The suitable ligand for this complex is :

- Options
1. ethylenediamine
 2. CN^-
 3. NCS^-
 4. CO

Question ID : 4165299913
Option 1 ID : 41652939113
Option 2 ID : 41652939112
Option 3 ID : 41652939111
Option 4 ID : 41652939110
Status : Not Answered
Chosen Option : --

Q.3 The element that does NOT show catenation is :

- Options
1. Ge
 2. Si
 3. Sn
 4. Pb

Question ID : 4165299910
Option 1 ID : 41652939099
Option 2 ID : 41652939098
Option 3 ID : 41652939100
Option 4 ID : 41652939101
Status : Answered
Chosen Option : 2

Q.4 Among the following, the false statement is :

- Options
1. It is possible to cause artificial rain by throwing electrified sand carrying charge opposite to the one on clouds from an aeroplane.
 2. Tyndall effect can be used to distinguish between a colloidal solution and a true solution.

3. Lyophilic sol can be coagulated by adding an electrolyte.

4. Latex is a colloidal solution of rubber particles which are positively charged

Question ID : 4165299925

Option 1 ID : 41652939160

Option 2 ID : 41652939158

Option 3 ID : 41652939161





Option 4 ID : 41652939159

Status : Answered

Chosen Option : 1

Q.5 The correct structure of histidine in a strongly acidic solution (pH = 2) is :

Options



Question ID : 4165299899

Option 1 ID : 41652939056

Option 2 ID : 41652939057

Option 3 ID : 41652939055


Option 4 ID : 41652939054

Status : Answered

Chosen Option : 1

Q.6

The major product of the following reaction is :



(i) NaNO_2/H^+

(ii) CrO_3/H^+

(iii) H_2SO_4 (conc.), Δ

Options

- 1.
- 2.
- 3.
- 4.

Question ID : 4165299898
 Option 1 ID : 41652939051
 Option 2 ID : 41652939053
 Option 3 ID : 41652939052
 Option 4 ID : 41652939050
 Status : Answered
 Chosen Option : 2

Q.7

\wedge_m° for NaCl, HCl and NaA are 126.4, 425.9 and $100.5 \text{ S cm}^2\text{mol}^{-1}$, respectively. If the conductivity of 0.001 M HA is $5 \times 10^{-5} \text{ S cm}^{-1}$, degree of dissociation of HA is :

- Options
1. 0.50
 2. 0.25
 3. 0.125
 4. 0.75

Question ID : 4165299923
 Option 1 ID : 41652939152
 Option 2 ID : 41652939150
 Option 3 ID : 41652939151
 Option 4 ID : 41652939153
 Status : Answered

Chosen Option : 3

Q.8 Molecules of benzoic acid (C_6H_5COOH) dimerise in benzene. 'w' g of the acid dissolved in 30 g of benzene shows a depression in freezing point equal to 2 K. If the percentage association of the acid to form dimer in the solution is 80, then w is :
(Given that $K_f = 5 \text{ K kg mol}^{-1}$, Molar mass of benzoic acid = 122 g mol^{-1})

- Options
1. 2.4 g
 2. 1.0 g
 3. 1.5 g
 4. 1.8 g

Question ID : 4165299921

Option 1 ID : 41652939144

Option 2 ID : 41652939142

Option 3 ID : 41652939145

Option 4 ID : 41652939143

Status : Not Answered

Chosen Option : --

Q.9 Chlorine on reaction with hot and concentrated sodium hydroxide gives :

- Options
1. Cl^- and ClO_3^-
 2. Cl^- and ClO^-
 3. ClO_3^- and ClO_2^-
 4. Cl^- and ClO_2^-

Question ID : 4165299911

Option 1 ID : 41652939103

Option 2 ID : 41652939105


Option 3 ID : 41652939102

Option 4 ID : 41652939104


Status : Answered

Chosen Option : 4

Q.10 The major product of the following reaction is :



- Options
1. $\text{CH}_3\text{CH}_2\text{C}=\text{CH}_2$
|
 $\text{CO}_2\text{CH}_2\text{CH}_3$
 2. $\text{CO}_2\text{CH}_2\text{CH}_3$
|
 $\text{CH}_3\text{C}=\text{CHCH}_3$



Question ID : 4165299905

Option 1 ID : 41652939081

Option 2 ID : 41652939079


Option 3 ID : 41652939078

Option 4 ID : 41652939080

Status : Not Answered

Chosen Option : --

Q.11 The combination of plots which does not represent isothermal expansion of an ideal gas is :



- Options
1. (B) and (D)
 2. (A) and (C)
 3. (B) and (C)
 4. (A) and (D)

Question ID : 4165299919

Option 1 ID : 41652939136

Option 2 ID : 41652939137

Option 3 ID : 41652939134


Option 4 ID : 41652939135

Status : Answered





Chosen Option : 1

Q.12

The major product in the following conversion is :



Options

1. 
2. 
3. 
4. 

Question ID : 4165299904

Option 1 ID : 41652939076

Option 2 ID : 41652939077


Option 3 ID : 41652939074

Option 4 ID : 41652939075





Status : Answered

Chosen Option : 4

Q.13 The major product of the following reaction is :



Options

1. 
2. 
3. 
4. 

Question ID : 4165299900

Option 1 ID : 41652939061

Option 2 ID : 41652939059

Option 3 ID : **41652939060**
 Option 4 ID : **41652939058**
 Status : **Answered**
 Chosen Option : **1**

Q.14 The correct order of atomic radii is :

- Options
1. N > Ce > Eu > Ho
 2. Ho > N > Eu > Ce
 3. Ce > Eu > Ho > N
 4. Eu > Ce > Ho > N

Question ID : **4165299912**
 Option 1 ID : **41652939108**
 Option 2 ID : **41652939109**
 Option 3 ID : **41652939107**
 Option 4 ID : **41652939106**
 Status : **Answered**
 Chosen Option : **3**

Q.15 The element that shows greater ability to form $p\pi - p\pi$ multiple bonds, is :

- Options
1. Sn
 2. C
 3. Ge
 4. Si

Question ID : **4165299906**
 Option 1 ID : **41652939085**
 Option 2 ID : **41652939083**
 Option 3 ID : **41652939084**
 Option 4 ID : **41652939082**
 Status : **Answered**
 Chosen Option : **2**





Q.16 The two monomers for the synthesis of Nylon 6, 6 are :

- Options
1. $\text{HOOC}(\text{CH}_2)_4\text{COOH}$,
 $\text{H}_2\text{N}(\text{CH}_2)_6\text{NH}_2$
 2. $\text{HOOC}(\text{CH}_2)_6\text{COOH}$,
 $\text{H}_2\text{N}(\text{CH}_2)_6\text{NH}_2$
 3. $\text{HOOC}(\text{CH}_2)_4\text{COOH}$,
 $\text{H}_2\text{N}(\text{CH}_2)_4\text{NH}_2$
 4. $\text{HOOC}(\text{CH}_2)_6\text{COOH}$,
 $\text{H}_2\text{N}(\text{CH}_2)_4\text{NH}_2$

Question ID : **4165299897**
 Option 1 ID : **41652939048**
 Option 2 ID : **41652939046**
 Option 3 ID : **41652939047**
 Option 4 ID : **41652939049**
 Status : **Not Answered**
 Chosen Option : **--**

Q.17

The aldehydes which will **not** form Grignard product with one equivalent Grignard reagents are :

- (A) 
- (B) 
- (C) 
- (D) 

- Options
1. (B), (D)
 2. (B), (C)
 3. (B), (C), (D)
 4. (C), (D)

Question ID : 4165299901

Option 1 ID : 41652939063

Option 2 ID : 41652939062




Option 3 ID : 41652939064

Option 4 ID : 41652939065

Status : Not Answered

Chosen Option : --

Q.18 Given :



Based on the above thermochemical equations, find out which one of the following algebraic relationships is correct ?

- Options
1. $x = y + z$
 2. $z = x + y$
 3. $y = 2z - x$
 4. $x = y - z$

Question ID : 4165299920

Option 1 ID : 41652939138

Option 2 ID : 41652939140

Option 3 ID : 41652939141

Option 4 ID : **41652939139**
 Status : **Not Answered**
 Chosen Option : --

Q.19 The volume strength of 1M H_2O_2 is :
 (Molar mass of $\text{H}_2\text{O}_2 = 34 \text{ g mol}^{-1}$)

- Options
1. 5.6
 2. 16.8
 3. 11.35
 4. 22.4

Question ID : **4165299908**
 Option 1 ID : **41652939090**
 Option 2 ID : **41652939092**
 Option 3 ID : **41652939091**
 Option 4 ID : **41652939093**
 Status : **Answered**
 Chosen Option : **4**

Q.20 The correct statement(s) among I to III with respect to potassium ions that are abundant within the cell fluids is/are :

- I. They activate many enzymes
- II. They participate in the oxidation of glucose to produce ATP
- III. Along with sodium ions, they are responsible for the transmission of nerve signals

- Options
1. I and II only
 2. I and III only
 3. I, II and III
 4. III only

Question ID : **4165299909**
 Option 1 ID : **41652939095**
 Option 2 ID : **41652939097**
 Option 3 ID : **41652939094**
 Option 4 ID : **41652939096**
 Status : **Answered**
 Chosen Option : **2**

Q.21 The compound that is NOT a common component of photochemical smog is :

- Options
1. O_3
 2. $\text{H}_3\text{C}-\overset{\text{O}}{\parallel}{\text{C}}-\text{OONO}_2$
 3. $\text{CH}_2=\text{CHCHO}$
 4. CF_2Cl_2


Question ID : **4165299915**
 Option 1 ID : **41652939118**
 Option 2 ID : **41652939121**
 Option 3 ID : **41652939119**

Option 4 ID : 41652939120

Status : Answered

Chosen Option : 3

- Q.22** For a reaction, consider the plot of $\ln k$ versus $1/T$ given in the figure. If the rate constant of this reaction at 400 K is 10^{-5} s^{-1} , then the rate constant at 500 K is :



- Options**
1. 10^{-6} s^{-1}
 2. $2 \times 10^{-4} \text{ s}^{-1}$
 3. 10^{-4} s^{-1}
 4. $4 \times 10^{-4} \text{ s}^{-1}$

Question ID : 4165299924

Option 1 ID : 41652939155

Option 2 ID : 41652939156

Option 3 ID : 41652939154

Option 4 ID : 41652939157

Status : Answered

Chosen Option : 3

- Q.23** An open vessel at 27°C is heated until two fifth of the air (assumed as an ideal gas) in it has escaped from the vessel. Assuming that the volume of the vessel remains constant, the temperature at which the vessel has been heated is :

- Options**
1. 500°C
 2. 500 K
 3. 750°C
 4. 750 K

Question ID : 4165299917

Option 1 ID : 41652939126

Option 2 ID : 41652939127


Option 3 ID : 41652939129

Option 4 ID : 41652939128


Status : Answered

Chosen Option : 4

- Q.24** The major product of the following reaction is :



Options



Question ID : 4165299903

Option 1 ID : 41652939073

Option 2 ID : 41652939070


Option 3 ID : 41652939071

Option 4 ID : 41652939072

Status : Not Answered

Chosen Option : --

Q.25 The increasing order of the reactivity of the following with LiAlH_4 is :



- Options
1. (B) < (A) < (C) < (D)
 2. (B) < (A) < (D) < (C)
 3. (A) < (B) < (D) < (C)
 4. (A) < (B) < (C) < (D)

Question ID : 4165299902

Option 1 ID : 41652939069

Option 2 ID : 41652939067

Option 3 ID : 41652939068

Option 4 ID : 41652939066

Status : Answered

Chosen Option : 4

Q.26

The pair that does NOT require calcination is :

- Options
1. ZnO and MgO
 2. ZnO and $\text{Fe}_2\text{O}_3 \cdot x\text{H}_2\text{O}$
 3. ZnCO_3 and CaO
 4. Fe_2O_3 and $\text{CaCO}_3 \cdot \text{MgCO}_3$

Question ID : 4165299907

Option 1 ID : 41652939088

Option 2 ID : 41652939086


Option 3 ID : 41652939087

Option 4 ID : 41652939089

Status : Answered

Chosen Option : 1

Q.27 The major product of the following reaction is :



- Options
1. $\text{CH}_3\text{CH}=\text{C}=\text{CH}_2$
 2. $\text{CH}_3\text{CH}_2\underset{\text{NH}_2}{\text{CH}}-\underset{\text{NH}_2}{\text{CH}_2}$
 3. $\text{CH}_3\text{CH}=\text{CHCH}_2\text{NH}_2$
 4. $\text{CH}_3\text{CH}_2\text{C}\equiv\text{CH}$

Question ID : 4165299896

Option 1 ID : 41652939043

Option 2 ID : 41652939044

Option 3 ID : 41652939042

Option 4 ID : 41652939045

Status : Answered

Chosen Option : 3

Q.28 If K_{sp} of Ag_2CO_3 is 8×10^{-12} , the molar solubility of Ag_2CO_3 in 0.1 M AgNO_3 is :

- Options
1. 8×10^{-12} M
 2. 8×10^{-11} M
 3. 8×10^{-10} M
 4. 8×10^{-13} M

Question ID : 4165299922

Option 1 ID : 41652939148

Option 2 ID : 41652939147

Option 3 ID : 41652939146

Option 4 ID : 41652939149

Status : Not Answered

Chosen Option : --

Q.29

The upper stratosphere consisting of the ozone layer protects us from the sun's radiation that falls in the wavelength region of :

- Options
1. 200 - 315 nm
 2. 400 - 550 nm
 3. 0.8 - 1.5 nm
 4. 600 - 750 nm

Question ID : 4165299914
 Option 1 ID : 41652939115
 Option 2 ID : 41652939116
 Option 3 ID : 41652939114
 Option 4 ID : 41652939117
 Status : Answered
 Chosen Option : 2

Q.30 If the de Broglie wavelength of the electron in n^{th} Bohr orbit in a hydrogenic atom is equal to $1.5 \pi a_0$ (a_0 is Bohr radius), then the value of n/z is :

- Options
1. 0.40
 2. 1.50
 3. 1.0
 4. 0.75

Question ID : 4165299918
 Option 1 ID : 41652939133
 Option 2 ID : 41652939131
 Option 3 ID : 41652939130
 Option 4 ID : 41652939132
 Status : Answered
 Chosen Option : 2

Section : Mathematics

Q.1

$\lim_{x \rightarrow 1^-} \frac{\sqrt{\pi} - \sqrt{2 \sin^{-1} x}}{\sqrt{1-x}}$ is equal to :

- Options
1. $\frac{1}{\sqrt{2\pi}}$
 2. $\sqrt{\frac{2}{\pi}}$
 3. $\sqrt{\frac{\pi}{2}}$
 4. $\sqrt{\pi}$

Question ID : 4165299935
 Option 1 ID : 41652939200
 Option 2 ID : 41652939198
 Option 3 ID : 41652939199

Option 4 ID : **41652939201**
 Status : **Answered**
 Chosen Option : **2**

Q.2 Let f be a differentiable function such that $f(1) = 2$ and $f'(x) = f(x)$ for all $x \in \mathbb{R}$. If $h(x) = f(f(x))$, then $h'(1)$ is equal to :

- Options
1. $2e^2$
 2. $4e$
 3. $2e$
 4. $4e^2$

Question ID : **4165299936**
 Option 1 ID : **41652939202**
 Option 2 ID : **41652939203**
 Option 3 ID : **41652939204**
 Option 4 ID : **41652939205**
 Status : **Not Answered**
 Chosen Option : --

Q.3

The integral $\int_1^e \left\{ \left(\frac{x}{e} \right)^{2x} - \left(\frac{e}{x} \right)^x \right\} \log_e x \, dx$

is equal to :

- Options
1. $\frac{1}{2} - e - \frac{1}{e^2}$
 2. $-\frac{1}{2} + \frac{1}{e} - \frac{1}{2e^2}$
 3. $\frac{3}{2} - \frac{1}{e} - \frac{1}{2e^2}$
 4. $\frac{3}{2} - e - \frac{1}{2e^2}$

Question ID : **4165299940**
 Option 1 ID : **41652939221**
 Option 2 ID : **41652939220**
 Option 3 ID : **41652939218**
 Option 4 ID : **41652939219**
 Status : **Not Answered**
 Chosen Option : --

Q.4

Let \vec{a} , \vec{b} and \vec{c} be three unit vectors, out of which vectors \vec{b} and \vec{c} are non-parallel. If α and β are the angles which vector \vec{a} makes with vectors \vec{b} and \vec{c} respectively and $\vec{a} \times (\vec{b} \times \vec{c}) = \frac{1}{2} \vec{b}$, then $|\alpha - \beta|$ is equal to :

- Options
1. 30°
 2. 90°
 3. 60°
 4. 45°

Question ID : 4165299949

Option 1 ID : 41652939256

Option 2 ID : 41652939257

Option 3 ID : 41652939254

Option 4 ID : 41652939255

Status : Not Answered

Chosen Option : --

Q.5

The integral $\int \frac{3x^{13} + 2x^{11}}{(2x^4 + 3x^2 + 1)^4} dx$ is

equal to :
(where C is a constant of integration)

Options

1. $\frac{x^4}{6(2x^4 + 3x^2 + 1)^3} + C$

2. $\frac{x^{12}}{6(2x^4 + 3x^2 + 1)^3} + C$

3. $\frac{x^4}{(2x^4 + 3x^2 + 1)^3} + C$

4. $\frac{x^{12}}{(2x^4 + 3x^2 + 1)^3} + C$

Question ID : 4165299939

Option 1 ID : 41652939214

Option 2 ID : 41652939215

Option 3 ID : 41652939216

Option 4 ID : 41652939217

Status : Not Answered

Chosen Option : --

Q.6 If $\sin^4 \alpha + 4 \cos^4 \beta + 2 = 4\sqrt{2} \sin \alpha \cos \beta$;
 $\alpha, \beta \in [0, \pi]$, then $\cos(\alpha + \beta) - \cos(\alpha - \beta)$ is
 equal to :

- Options
1. 0
 2. -1
 3. $\sqrt{2}$
 4. $-\sqrt{2}$

Question ID : 4165299953
 Option 1 ID : 41652939272
 Option 2 ID : 41652939273
 Option 3 ID : 41652939270
 Option 4 ID : 41652939271
 Status : Not Answered
 Chosen Option : --

Q.7 If a curve passes through the point $(1, -2)$
 and has slope of the tangent at any point
 (x, y) on it as $\frac{x^2 - 2y}{x}$, then the curve also
 passes through the point :

- Options
1. $(3, 0)$
 2. $(\sqrt{3}, 0)$
 3. $(-1, 2)$
 4. $(-\sqrt{2}, 1)$

Question ID : 4165299942
 Option 1 ID : 41652939226
 Option 2 ID : 41652939229
 Option 3 ID : 41652939227
 Option 4 ID : 41652939228
 Status : Not Answered
 Chosen Option : --

Q.8 If an angle between the line,
 $\frac{x+1}{2} = \frac{y-2}{1} = \frac{z-3}{-2}$ and the plane,
 $x - 2y - kz = 3$ is $\cos^{-1}\left(\frac{2\sqrt{2}}{3}\right)$, then a value
 of k is :

- Options
1. $\sqrt{\frac{5}{3}}$
 2. $\sqrt{\frac{3}{5}}$
 3. $-\frac{3}{5}$

4. $-\frac{5}{3}$

Question ID : 4165299947
 Option 1 ID : 41652939246
 Option 2 ID : 41652939247
 Option 3 ID : 41652939248
 Option 4 ID : 41652939249
 Status : Answered
 Chosen Option : 2

Q.9 $\lim_{n \rightarrow \infty} \left(\frac{n}{n^2 + 1^2} + \frac{n}{n^2 + 2^2} + \frac{n}{n^2 + 3^2} + \dots + \frac{1}{5n} \right)$
 is equal to :

- Options
1. $\frac{\pi}{4}$
 2. $\tan^{-1}(3)$
 3. $\frac{\pi}{2}$
 4. $\tan^{-1}(2)$

Question ID : 4165299941
 Option 1 ID : 41652939224
 Option 2 ID : 41652939223
 Option 3 ID : 41652939225
 Option 4 ID : 41652939222
 Status : Not Answered
 Chosen Option : --

Q.10 In a game, a man wins Rs. 100 if he gets 5 or 6 on a throw of a fair die and loses Rs. 50 for getting any other number on the die. If he decides to throw the die either till he gets a five or a six or to a maximum of three throws, then his expected gain/loss (in rupees) is :

- Options
1. $\frac{400}{9}$ loss
 2. 0
 3. $\frac{400}{3}$ gain
 4. $\frac{400}{3}$ loss

Question ID : 4165299952
 Option 1 ID : 41652939266
 Option 2 ID : 41652939269
 Option 3 ID : 41652939268
 Option 4 ID : 41652939267
 Status : Answered
 Chosen Option : 4

Q.11

If a straight line passing through the point $P(-3, 4)$ is such that its intercepted portion between the coordinate axes is bisected at P , then its equation is :

- Options
1. $3x - 4y + 25 = 0$
 2. $4x - 3y + 24 = 0$
 3. $x - y + 7 = 0$
 4. $4x + 3y = 0$

Question ID : 4165299943

Option 1 ID : 41652939232

Option 2 ID : 41652939233

Option 3 ID : 41652939231

Option 4 ID : 41652939230

Status : Answered

Chosen Option : 2

Q.12

There are m men and two women participating in a chess tournament. Each participant plays two games with every other participant. If the number of games played by the men between themselves exceeds the number of games played between the men and the women by 84, then the value of m is :

- Options
1. 12
 2. 11
 3. 9
 4. 7

Question ID : 4165299931

Option 1 ID : 41652939185

Option 2 ID : 41652939184

Option 3 ID : 41652939183

Option 4 ID : 41652939182

Status : Not Answered

Chosen Option : --

Q.13

The tangent to the curve $y = x^2 - 5x + 5$, parallel to the line $2y = 4x + 1$, also passes through the point :

- Options
1. $\left(\frac{7}{2}, \frac{1}{4}\right)$
 2. $\left(\frac{1}{8}, -7\right)$
 3. $\left(-\frac{1}{8}, 7\right)$
 4. $\left(\frac{1}{4}, \frac{7}{2}\right)$

Question ID : 4165299937
 Option 1 ID : 41652939206
 Option 2 ID : 41652939209
 Option 3 ID : 41652939208
 Option 4 ID : 41652939207
 Status : Answered
 Chosen Option : 4

Q.14 Let S be the set of all real values of λ such that a plane passing through the points $(-\lambda^2, 1, 1)$, $(1, -\lambda^2, 1)$ and $(1, 1, -\lambda^2)$ also passes through the point $(-1, -1, 1)$. Then S is equal to :

- Options
1. $\{\sqrt{3}\}$
 2. $\{\sqrt{3}, -\sqrt{3}\}$
 3. $\{1, -1\}$
 4. $\{3, -3\}$

Question ID : 4165299948
 Option 1 ID : 41652939250
 Option 2 ID : 41652939253
 Option 3 ID : 41652939251
 Option 4 ID : 41652939252
 Status : Answered
 Chosen Option : 2

Q.15 Let z_1 and z_2 be two complex numbers satisfying $|z_1| = 9$ and $|z_2 - 3 - 4i| = 4$. Then the minimum value of $|z_1 - z_2|$ is :

- Options
1. 0
 2. $\sqrt{2}$
 3. 1
 4. 2

Question ID : 4165299928
 Option 1 ID : 41652939170
 Option 2 ID : 41652939172
 Option 3 ID : 41652939171
 Option 4 ID : 41652939173
 Status : Answered
 Chosen Option : 1

Q.16 The total number of irrational terms in the binomial expansion of $\left(7^{1/5} - 3^{1/10}\right)^{60}$ is :

- Options
1. 55
 2. 49
 3. 48
 4. 54

Question ID : 4165299932

Option 1 ID : 41652939189

Option 2 ID : 41652939187

Option 3 ID : 41652939186

Option 4 ID : 41652939188

Status : Answered

Chosen Option : 3

Q.17 The equation of a tangent to the parabola, $x^2=8y$, which makes an angle θ with the positive direction of x -axis, is :

- Options
1. $y = x \tan\theta + 2 \cot\theta$
 2. $y = x \tan\theta - 2 \cot\theta$
 3. $x = y \cot\theta + 2 \tan\theta$
 4. $x = y \cot\theta - 2 \tan\theta$

Question ID : 4165299945

Option 1 ID : 41652939241

Option 2 ID : 41652939238

Option 3 ID : 41652939239

Option 4 ID : 41652939240

Status : Not Answered

Chosen Option : --

Q.18 In a class of 60 students, 40 opted for NCC, 30 opted for NSS and 20 opted for both NCC and NSS. If one of these students is selected at random, then the probability that the student selected has opted neither for NCC nor for NSS is :

- Options
1. $\frac{1}{6}$
 2. $\frac{1}{3}$
 3. $\frac{2}{3}$
 4. $\frac{5}{6}$

Question ID : 4165299951

Option 1 ID : 41652939262

Option 2 ID : 41652939265

Option 3 ID : 41652939263

Option 4 ID : 41652939264

Status : Answered

Chosen Option : 3

Q.19 The mean and the variance of five observations are 4 and 5.20, respectively. If three of the observations are 3, 4 and 4 ; then the absolute value of the difference of the other two observations, is :

- Options
1. 7
 2. 5

3. 1

4. 3

Question ID : 4165299950

Option 1 ID : 41652939261

Option 2 ID : 41652939260

Option 3 ID : 41652939258

Option 4 ID : 41652939259

Status : Answered

Chosen Option : 1

Q.20

If $A = \begin{bmatrix} 1 & \sin\theta & 1 \\ -\sin\theta & 1 & \sin\theta \\ -1 & -\sin\theta & 1 \end{bmatrix}$; then for all

$\theta \in \left(\frac{3\pi}{4}, \frac{5\pi}{4}\right)$, $\det(A)$ lies in the interval :

Options

1. $\left(1, \frac{5}{2}\right]$

2. $\left[\frac{5}{2}, 4\right)$

3. $\left(0, \frac{3}{2}\right]$

4. $\left(\frac{3}{2}, 3\right]$

Question ID : 4165299929

Option 1 ID : 41652939175

Option 2 ID : 41652939177

Option 3 ID : 41652939174

Option 4 ID : 41652939176

Status : Answered

Chosen Option : 3

Q.21

If a circle of radius R passes through the origin O and intersects the coordinate axes at A and B , then the locus of the foot of perpendicular from O on AB is :

Options 1. $(x^2 + y^2)^2 = 4R^2x^2y^2$

2. $(x^2 + y^2)^3 = 4R^2x^2y^2$

3. $(x^2 + y^2)^2 = 4R^2x^2y^2$

4. $(x^2 + y^2)(x + y) = R^2xy$

Question ID : 4165299944

Option 1 ID : 41652939235

Option 2 ID : 41652939234

Option 3 ID : 41652939236

Option 4 ID : 41652939237

Status : Answered

Chosen Option : 1

Q.22 The set of all values of λ for which the system of linear equations

$$x - 2y - 2z = \lambda x$$

$$x + 2y + z = \lambda y$$

$$-x - y = \lambda z$$

has a non-trivial solution :

- Options**
1. is a singleton
 2. contains exactly two elements
 3. is an empty set
 4. contains more than two elements

Question ID : 4165299930

Option 1 ID : 41652939178

Option 2 ID : 41652939180

Option 3 ID : 41652939179

Option 4 ID : 41652939181

Status : Answered

Chosen Option : 2

Q.23 If the function f given by

$f(x) = x^3 - 3(a-2)x^2 + 3ax + 7$, for some $a \in \mathbb{R}$ is increasing in $(0, 1]$ and decreasing in $[1, 5)$, then a root of the equation,

$$\frac{f(x) - 14}{(x - 1)^2} = 0 \quad (x \neq 1) \text{ is :}$$

- Options**
1. -7
 2. 5
 3. 7
 4. 6

Question ID : 4165299938

Option 1 ID : 41652939210

Option 2 ID : 41652939211

Option 3 ID : 41652939213

Option 4 ID : 41652939212

Status : Answered

Chosen Option : 3

Q.24 Let Z be the set of integers. If $A = \{x \in Z : 2(x+2)(x^2 - 5x + 6) = 1\}$ and $B = \{x \in Z : -3 < 2x - 1 < 9\}$, then the number of subsets of the set $A \times B$, is :

- Options**
1. 2^{15}
 2. 2^{18}
 3. 2^{12}
 4. 2^{10}

Question ID : 4165299926

Option 1 ID : 41652939164

Option 2 ID : 41652939165

Option 3 ID : 41652939163

Option 4 ID : 41652939162

Status : **Not Answered**
Chosen Option : --

Q.25 If nC_4 , nC_5 and nC_6 are in A.P., then n can be :

- Options
1. 9
 2. 14
 3. 11
 4. 12

Question ID : **4165299933**
Option 1 ID : **41652939190**
Option 2 ID : **41652939193**
Option 3 ID : **41652939191**
Option 4 ID : **41652939192**
Status : **Answered**
Chosen Option : **2**

Q.26 Let S and S' be the foci of an ellipse and B be any one of the extremities of its minor axis. If $\Delta S'BS$ is a right angled triangle with right angle at B and area $(\Delta S'BS) = 8$ sq. units, then the length of a latus rectum of the ellipse is :

- Options
1. 4
 2. $2\sqrt{2}$
 3. $4\sqrt{2}$
 4. 2

Question ID : **4165299946**
Option 1 ID : **41652939245**
Option 2 ID : **41652939243**
Option 3 ID : **41652939244**
Option 4 ID : **41652939242**
Status : **Not Answered**
Chosen Option : --

Q.27 If the angle of elevation of a cloud from a point P which is 25 m above a lake be 30° and the angle of depression of reflection of the cloud in the lake from P be 60° , then the height of the cloud (in meters) from the surface of the lake is :

- Options
1. 60
 2. 50
 3. 45
 4. 42

Question ID : **4165299954**
Option 1 ID : **41652939274**
Option 2 ID : **41652939275**
Option 3 ID : **41652939276**
Option 4 ID : **41652939277**
Status : **Answered**

Chosen Option : 4

Q.28 The expression $\sim(\sim p \rightarrow q)$ is logically equivalent to :

- Options
1. $\sim p \wedge \sim q$
 2. $p \wedge \sim q$
 3. $\sim p \wedge q$
 4. $p \wedge q$

Question ID : 4165299955

Option 1 ID : 41652939281

Option 2 ID : 41652939279

Option 3 ID : 41652939280

Option 4 ID : 41652939278

Status : Answered

Chosen Option : 4

Q.29 If the sum of the first 15 terms of the series

$$\left(\frac{3}{4}\right)^3 + \left(1\frac{1}{2}\right)^3 + \left(2\frac{1}{4}\right)^3 + 3^3 + \left(3\frac{3}{4}\right)^3 + \dots$$

is equal to 225 k, then k is equal to :

- Options
1. 108
 2. 27
 3. 54
 4. 9

Question ID : 4165299934

Option 1 ID : 41652939197

Option 2 ID : 41652939195

Option 3 ID : 41652939196

Option 4 ID : 41652939194

Status : Not Answered

Chosen Option : --

Q.30 The number of integral values of m for which the quadratic expression, $(1 + 2m)x^2 - 2(1 + 3m)x + 4(1 + m)$, $x \in \mathbb{R}$, is always positive, is :

- Options
1. 3
 2. 8
 3. 7
 4. 6

Question ID : 4165299927

Option 1 ID : 41652939169

Option 2 ID : 41652939166

Option 3 ID : 41652939167

Option 4 ID : 41652939168

Status : Not Answered

Chosen Option : --