

**JEE MAIN 2019**

Application No	
Candidate Name	
Roll No.	
Test Date	09/01/2019
Test Time	2:30 PM - 5:30 PM
Subject	Paper I EH

## Section : Physics

**Q.1** At a given instant, say  $t = 0$ , two radioactive substances A and B have equal activities.

The ratio  $\frac{R_B}{R_A}$  of their activities after

time  $t$  itself decays with time  $t$  as  $e^{-3t}$ .  
If the half-life of A is  $\ln 2$ , the half-life of B is :

- Options
1.  $4\ln 2$
  2.  $\frac{\ln 2}{2}$
  3.  $\frac{\ln 2}{4}$
  4.  $2\ln 2$

Question ID : 4165298811

Option 1 ID : 41652934705

Option 2 ID : 41652934703

Option 3 ID : 41652934702

Option 4 ID : 41652934704

Status : Marked For Review

Chosen Option : 2

**Q.2** A power transmission line feeds input power at 2300 V to a step down transformer with its primary windings having 4000 turns. The output power is delivered at 230 V by the transformer. If the current in the primary of the transformer is 5A and its efficiency is 90%, the output current would be :

- Options
1. 50 A
  2. 45 A
  3. 35 A
  4. 25 A

Question ID : 4165298806

Option 1 ID : 41652934683

Option 2 ID : 41652934684

Option 3 ID : 41652934682

Option 4 ID : 41652934685

Status : Marked For Review

Chosen Option : 2

Q.3 The energy associated with electric field is ( $U_E$ ) and with magnetic field is ( $U_B$ ) for an electromagnetic wave in free space. Then :

Options

1.  $U_E = \frac{U_B}{2}$
2.  $U_E > U_B$
3.  $U_E < U_B$
4.  $U_E = U_B$

Question ID : 4165298807  
Option 1 ID : 41652934689  
Option 2 ID : 41652934686  
Option 3 ID : 41652934687  
Option 4 ID : 41652934688

Status : Answered

Chosen Option : 3

Q.4 A force acts on a 2 kg object so that its position is given as a function of time as  $x = 3t^2 + 5$ . What is the work done by this force in first 5 seconds ?

Options

1. 850 J
2. 950 J
3. 875 J
4. 900 J

Question ID : 4165298790  
Option 1 ID : 41652934621  
Option 2 ID : 41652934618  
Option 3 ID : 41652934620  
Option 4 ID : 41652934619

Status : Answered

Chosen Option : 3

Q.5 A particle having the same charge as of electron moves in a circular path of radius 0.5 cm under the influence of a magnetic field of 0.5 T. If an electric field of 100 V/m makes it to move in a straight path, then the mass of the particle is (Given charge of electron =  $1.6 \times 10^{-19} \text{C}$ )

Options

1.  $9.1 \times 10^{-31} \text{ kg}$
2.  $1.6 \times 10^{-27} \text{ kg}$
3.  $1.6 \times 10^{-19} \text{ kg}$
4.  $2.0 \times 10^{-24} \text{ kg}$

Question ID : 4165298805  
Option 1 ID : 41652934678  
Option 2 ID : 41652934681  
Option 3 ID : 41652934680

Option 4 ID : 41652934679  
 Status : Answered  
 Chosen Option : 1

Q.6 Two point charges  $q_1(\sqrt{10} \mu\text{C})$  and  $q_2(-25 \mu\text{C})$  are placed on the  $x$ -axis at  $x=1 \text{ m}$  and  $x=4 \text{ m}$  respectively. The electric field (in  $\text{V/m}$ ) at a point  $y=3 \text{ m}$  on  $y$ -axis is,

$$\left[ \text{take } \frac{1}{4\pi\epsilon_0} = 9 \times 10^9 \text{ Nm}^2\text{C}^{-2} \right]$$

Options

1.  $(63 \hat{i} - 27 \hat{j}) \times 10^2$
2.  $(-63 \hat{i} + 27 \hat{j}) \times 10^2$
3.  $(81 \hat{i} - 81 \hat{j}) \times 10^2$
4.  $(-81 \hat{i} + 81 \hat{j}) \times 10^2$

Question ID : 4165298801  
 Option 1 ID : 41652934663  
 Option 2 ID : 41652934662  
 Option 3 ID : 41652934664  
 Option 4 ID : 41652934665  
 Status : Not Answered  
 Chosen Option : --

Q.7 Expression for time in terms of  $G$  (universal gravitational constant),  $h$  (Planck constant) and  $c$  (speed of light) is proportional to :


Options

1.  $\sqrt{\frac{hc^5}{G}}$
2.  $\sqrt{\frac{c^3}{Gh}}$
3.  $\sqrt{\frac{Gh}{c^5}}$
4.  $\sqrt{\frac{Gh}{c^3}}$

Question ID : 4165298786  
 Option 1 ID : 41652934605  
 Option 2 ID : 41652934603  
 Option 3 ID : 41652934604  
 Option 4 ID : 41652934602  
 Status : Answered  
 Chosen Option : 4

Q.8

Ge and Si diodes start conducting at 0.3 V and 0.7 V respectively. In the following figure if Ge diode connection are reversed, the value of  $V_o$  changes by : (assume that the Ge diode has large breakdown voltage)



- Options
1. 0.8 V
  2. 0.6 V
  3. 0.2 V
  4. 0.4 V

Question ID : 4165298812  
 Option 1 ID : 41652934709  
 Option 2 ID : 41652934708  
 Option 3 ID : 41652934707  
 Option 4 ID : 41652934706  
 Status : Not Answered

Chosen Option : --

Q.9 The top of a water tank is open to air and its water level is maintained. It is giving out  $0.74 \text{ m}^3$  water per minute through a circular opening of 2 cm radius in its wall. The depth of the centre of the opening from the level of water in the tank is close to :

- Options
1. 6.0 m
  2. 4.8 m
  3. 9.6 m
  4. 2.9 m

Question ID : 4165298794  
 Option 1 ID : 41652934637  
 Option 2 ID : 41652934634  
 Option 3 ID : 41652934635  
 Option 4 ID : 41652934636  
 Status : Not Answered

Chosen Option : --

Q.10 The energy required to take a satellite to a height 'h' above Earth surface (radius of Earth =  $6.4 \times 10^3$  km) is  $E_1$  and kinetic energy required for the satellite to be in a circular orbit at this height is  $E_2$ . The value of h for which  $E_1$  and  $E_2$  are equal, is :

- Options
1.  $1.6 \times 10^3$  km

2.  $3.2 \times 10^3$  km
3.  $6.4 \times 10^3$  km
4.  $1.28 \times 10^4$  km

Question ID : 4165298793

Option 1 ID : 41652934632

Option 2 ID : 41652934633

Option 3 ID : 41652934631

Option 4 ID : 41652934630

Status : Not Answered

Chosen Option : --

Q.11 Two Carnot engines A and B are operated in series. The first one, A, receives heat at  $T_1 (= 600 \text{ K})$  and rejects to a reservoir at temperature  $T_2$ . The second engine B receives heat rejected by the first engine and, in turn, rejects to a heat reservoir at  $T_3 (= 400 \text{ K})$ . Calculate the temperature  $T_2$  if the work outputs of the two engines are equal :

- Options
1. 600 K
  2. 400 K
  3. 300 K
  4. 500 K

Question ID : 4165298795

Option 1 ID : 41652934641

Option 2 ID : 41652934639

Option 3 ID : 41652934638

Option 4 ID : 41652934640

Status : Not Answered

Chosen Option : --

Q.12 A series AC circuit containing an inductor (20 mH), a capacitor (120  $\mu\text{F}$ ) and a resistor (60  $\Omega$ ) is driven by an AC source of 24 V/50 Hz. The energy dissipated in the circuit in 60 s is :

- Options
1.  $5.65 \times 10^2$  J
  2.  $2.26 \times 10^3$  J
  3.  $5.17 \times 10^2$  J
  4.  $3.39 \times 10^3$  J

Question ID : 4165298803

Option 1 ID : 41652934670

Option 2 ID : 41652934671

Option 3 ID : 41652934672

Option 4 ID : 41652934673

Status : Not Answered

Chosen Option : --

Q.13

A particle is executing simple harmonic motion (SHM) of amplitude  $A$ , along the  $x$ -axis, about  $x=0$ . When its potential Energy (PE) equals kinetic energy (KE), the position of the particle will be :

Options

1.  $\frac{A}{2}$
2.  $\frac{A}{2\sqrt{2}}$
3.  $\frac{A}{\sqrt{2}}$
4.  $A$

Question ID : 4165298798

Option 1 ID : 41652934651

Option 2 ID : 41652934653

Option 3 ID : 41652934652

Option 4 ID : 41652934650

Status : Not Answered

Chosen Option : --

Q.14 A mass of 10 kg is suspended vertically by a rope from the roof. When a horizontal force is applied on the rope at some point, the rope deviated at an angle of  $45^\circ$  at the roof point. If the suspended mass is at equilibrium, the magnitude of the force applied is ( $g = 10 \text{ ms}^{-2}$ )

Options

1. 200 N
2. 140 N
3. 70 N
4. 100 N

Question ID : 4165298789

Option 1 ID : 41652934615

Option 2 ID : 41652934616

Option 3 ID : 41652934617

Option 4 ID : 41652934614

Status : Not Answered

Chosen Option : --

Q.15 A 15 g mass of nitrogen gas is enclosed in a vessel at a temperature  $27^\circ\text{C}$ . Amount of heat transferred to the gas, so that rms velocity of molecules is doubled, is about :  
[Take  $R = 8.3 \text{ J/K mole}$ ]

Options

1. 0.9 kJ
2. 6 kJ
3. 10 kJ
4. 14 kJ

Question ID : 4165298796  
Option 1 ID : 41652934642  
Option 2 ID : 41652934643  
Option 3 ID : 41652934644  
Option 4 ID : 41652934645  
Status : Not Answered  
Chosen Option : --

Q.16 In a Young's double slit experiment, the slits are placed 0.320 mm apart. Light of wavelength  $\lambda = 500$  nm is incident on the slits. The total number of bright fringes that are observed in the angular range  $-30^\circ \leq \theta \leq 30^\circ$  is :

- Options
1. 640
  2. 320
  3. 321
  4. 641

Question ID : 4165298809  
Option 1 ID : 41652934696  
Option 2 ID : 41652934694  
Option 3 ID : 41652934695  
Option 4 ID : 41652934697  
Status : Not Answered  
Chosen Option : --


Q.17 Two plane mirrors are inclined to each other such that a ray of light incident on the first mirror ( $M_1$ ) and parallel to the second mirror ( $M_2$ ) is finally reflected from the second mirror ( $M_2$ ) parallel to the first mirror ( $M_1$ ). The angle between the two mirrors will be :

- Options
1.  $45^\circ$
  2.  $60^\circ$
  3.  $75^\circ$
  4.  $90^\circ$

Question ID : 4165298808  
Option 1 ID : 41652934690  
Option 2 ID : 41652934691  
Option 3 ID : 41652934692  
Option 4 ID : 41652934693  
Status : Answered  
Chosen Option : 4

Q.18

A rod of length 50 cm is pivoted at one end. It is raised such that it makes an angle of  $30^\circ$  from the horizontal as shown and released from rest. Its angular speed when it passes through the horizontal (in  $\text{rad s}^{-1}$ ) will be ( $g = 10 \text{ ms}^{-2}$ )



Options

1.  $\sqrt{\frac{30}{2}}$
2.  $\sqrt{30}$
3.  $\frac{\sqrt{20}}{3}$
4.  $\frac{\sqrt{30}}{2}$

Question ID : 4165298791

Option 1 ID : 41652934623

Option 2 ID : 41652934622

Option 3 ID : 41652934625


Option 4 ID : 41652934624

Status : Answered

Chosen Option : 2

Q.19

A carbon resistance has a following colour code. What is the value of the resistance ?



Options

1.  $530 \text{ k}\Omega \pm 5\%$
2.  $5.3 \text{ M}\Omega \pm 5\%$
3.  $6.4 \text{ M}\Omega \pm 5\%$
4.  $64 \text{ k}\Omega \pm 10\%$

Question ID : 4165298814

Option 1 ID : 41652934715

Option 2 ID : 41652934717

Option 3 ID : 41652934714

Option 4 ID : 41652934716

Status : Not Answered

Chosen Option : --

Q.20



One of the two identical conducting wires of length  $L$  is bent in the form of a circular loop and the other one into a circular coil of  $N$  identical turns. If the same current is passed in both, the ratio of the magnetic field at the central of the loop ( $B_L$ ) to that at

the centre of the coil ( $B_C$ ), i.e.  $\frac{B_L}{B_C}$  will be :

- Options
1.  $N$
  2.  $\frac{1}{N}$
  3.  $N^2$
  4.  $\frac{1}{N^2}$

Question ID : 4165298804

Option 1 ID : 41652934674

Option 2 ID : 41652934677

Option 3 ID : 41652934675

Option 4 ID : 41652934676

Status : Not Answered

Chosen Option : --

Q.21 A rod of mass ' $M$ ' and length ' $2L$ ' is suspended at its middle by a wire. It exhibits torsional oscillations; If two masses each of ' $m$ ' are attached at distance ' $L/2$ ' from its centre on both sides, it reduces the oscillation frequency by 20%. The value of ratio  $m/M$  is close to :

- Options
1. 0.77
  2. 0.57
  3. 0.37
  4. 0.17

Question ID : 4165298792

Option 1 ID : 41652934627

Option 2 ID : 41652934628

Option 3 ID : 41652934629

Option 4 ID : 41652934626

Status : Not Answered

Chosen Option : --

Q.22 Charge is distributed within a sphere of radius  $R$  with a volume charge density

$$\rho(r) = \frac{A}{r^2} e^{-2r/a}, \text{ where } A \text{ and } a \text{ are constants.}$$

If  $Q$  is the total charge of this charge distribution, the radius  $R$  is :

- Options
1.  $a \log \left( 1 - \frac{Q}{2\pi a A} \right)$

$$2. \frac{a}{2} \log \left( \frac{1}{1 - \frac{Q}{2\pi aA}} \right)$$

$$3. a \log \left( \frac{1}{1 - \frac{Q}{2\pi aA}} \right)$$

$$4. \frac{a}{2} \log \left( 1 - \frac{Q}{2\pi aA} \right)$$

Question ID : 4165298800

Option 1 ID : 41652934660

Option 2 ID : 41652934659


Option 3 ID : 41652934661

Option 4 ID : 41652934658

Status : Not Answered

Chosen Option : --

Q.23 A parallel plate capacitor with square plates is filled with four dielectrics of dielectric constants  $K_1, K_2, K_3, K_4$  arranged as shown in the figure. The effective dielectric constant  $K$  will be :



Options

$$1. K = \frac{(K_1 + K_3)(K_2 + K_4)}{K_1 + K_2 + K_3 + K_4}$$

$$2. K = \frac{(K_1 + K_2)(K_3 + K_4)}{2(K_1 + K_2 + K_3 + K_4)}$$

$$3. K = \frac{(K_1 + K_2)(K_3 + K_4)}{K_1 + K_2 + K_3 + K_4}$$

$$4. K = \frac{(K_1 + K_4)(K_2 + K_3)}{2(K_1 + K_2 + K_3 + K_4)}$$

Question ID : 4165298799

Option 1 ID : 41652934655

Option 2 ID : 41652934657

Option 3 ID : 41652934654

Option 4 ID : 41652934656

Status : Not Answered

Chosen Option : --

Q.24

The pitch and the number of divisions, on the circular scale, for a given screw gauge are 0.5 mm and 100 respectively. When the screw gauge is fully tightened without any object, the zero of its circular scale lies 3 divisions below the mean line.

The readings of the main scale and the circular scale, for a thin sheet, are 5.5 mm and 48 respectively, the thickness of this sheet is :

- Options
1. 5.755 mm
  2. 5.950 mm
  3. 5.725 mm
  4. 5.740 mm

Question ID : 4165298815

Option 1 ID : 41652934719

Option 2 ID : 41652934720

Option 3 ID : 41652934718

Option 4 ID : 41652934721

Status : Not Answered

Chosen Option : --

Q.25 A musician using an open flute of length 50 cm produces second harmonic sound waves. A person runs towards the musician from another end of a hall at a speed of 10 km/h. If the wave speed is 330 m/s, the frequency heard by the running person shall be close to :

- Options
1. 666 Hz
  2. 753 Hz
  3. 500 Hz
  4. 333 Hz

Question ID : 4165298797

Option 1 ID : 41652934648

Option 2 ID : 41652934649

Option 3 ID : 41652934647

Option 4 ID : 41652934646

Status : Not Answered

Chosen Option : --

Q.26 In a car race on straight road, car A takes a time  $t$  less than car B at the finish and passes finishing point with a speed ' $v$ ' more than that of car B. Both the cars start from rest and travel with constant acceleration  $a_1$  and  $a_2$  respectively. Then ' $v$ ' is equal to :

- Options
1.  $\frac{2a_1 a_2}{a_1 + a_2} t$

2.  $\sqrt{2a_1a_2} t$

3.  $\sqrt{a_1a_2} t$

4.  $\frac{a_1 + a_2}{2} t$

Question ID : 4165298787

Option 1 ID : 41652934609

Option 2 ID : 41652934606

Option 3 ID : 41652934607

Option 4 ID : 41652934608

Status : Not Answered

Chosen Option : --

Q.27

The magnetic field associated with a light wave is given, at the origin, by

$$B = B_0 [\sin(3.14 \times 10^7 ct) + \sin(6.28 \times 10^7 ct)].$$

If this light falls on a silver plate having a work function of 4.7 eV, what will be the maximum kinetic energy of the photo electrons ?

$$(c = 3 \times 10^8 \text{ ms}^{-1}, h = 6.6 \times 10^{-34} \text{ J-s})$$

Options 1. 6.82 eV

2. 12.5 eV

3. 8.52 eV

4. 7.72 eV

Question ID : 4165298810

Option 1 ID : 41652934701

Option 2 ID : 41652934700

Option 3 ID : 41652934699


Option 4 ID : 41652934698

Status : Not Answered

Chosen Option : --

Q.28

In the given circuit the internal resistance of the 18 V cell is negligible. If  $R_1 = 400 \Omega$ ,  $R_3 = 100 \Omega$  and  $R_4 = 500 \Omega$  and the reading of an ideal voltmeter across  $R_4$  is 5 V, then the value of  $R_2$  will be :

Options 1. 300  $\Omega$ 2. 450  $\Omega$ 3. 550  $\Omega$ 4. 230  $\Omega$

Question ID : 4165298802  
 Option 1 ID : 41652934669  
 Option 2 ID : 41652934668  
 Option 3 ID : 41652934666  
 Option 4 ID : 41652934667  
 Status : Answered  
 Chosen Option : 4

Q.29 In a communication system operating at wavelength 800 nm, only one percent of source frequency is available as signal bandwidth. The number of channels accommodated for transmitting TV signals of band width 6 MHz are (Take velocity of light  $c = 3 \times 10^8 \text{ m/s}$ ,  $h = 6.6 \times 10^{-34} \text{ J-s}$ )

- Options
1.  $3.75 \times 10^6$
  2.  $3.86 \times 10^6$
  3.  $6.25 \times 10^5$
  4.  $4.87 \times 10^5$

Question ID : 4165298813  
 Option 1 ID : 41652934713  
 Option 2 ID : 41652934711  
 Option 3 ID : 41652934710  
 Option 4 ID : 41652934712  
 Status : Not Answered  
 Chosen Option : --

Q.30 The position co-ordinates of a particle moving in a 3-D coordinate system is given by  $x = a \cos \omega t$   
 $y = a \sin \omega t$   
 and  $z = a \omega t$   
 The speed of the particle is :

- Options
1.  $\sqrt{2} a \omega$
  2.  $a \omega$
  3.  $\sqrt{3} a \omega$
  4.  $2a \omega$

Question ID : 4165298788  
 Option 1 ID : 41652934612  
 Option 2 ID : 41652934610  
 Option 3 ID : 41652934613  
 Option 4 ID : 41652934611  
 Status : Marked For Review  
 Chosen Option : 4

Section : Chemistry

Q.1

The entropy change associated with the conversion of 1 kg of ice at 273 K to water vapours at 383 K is :

(Specific heat of water liquid and water vapour are  $4.2 \text{ kJ K}^{-1} \text{ kg}^{-1}$  and  $2.0 \text{ kJ K}^{-1} \text{ kg}^{-1}$  ; heat of liquid fusion and vapourisation of water are  $334 \text{ kJ kg}^{-1}$  and  $2491 \text{ kJ kg}^{-1}$ , respectively). ( $\log 273 = 2.436$ ,  $\log 373 = 2.572$ ,  $\log 383 = 2.583$ )

- Options
1.  $7.90 \text{ kJ kg}^{-1} \text{ K}^{-1}$
  2.  $2.64 \text{ kJ kg}^{-1} \text{ K}^{-1}$
  3.  $8.49 \text{ kJ kg}^{-1} \text{ K}^{-1}$
  4.  $9.26 \text{ kJ kg}^{-1} \text{ K}^{-1}$

Question ID : 4165298840

Option 1 ID : 41652934820

Option 2 ID : 41652934818


Option 3 ID : 41652934819

Option 4 ID : 41652934821

Status : Not Answered

Chosen Option : --

Q.2 For the following reaction, the mass of water produced from 445 g of  $\text{C}_{57}\text{H}_{110}\text{O}_6$  is :



- Options
1. 490 g
  2. 445 g
  3. 495 g
  4. 890 g

Question ID : 4165298836

Option 1 ID : 41652934804

Option 2 ID : 41652934802


Option 3 ID : 41652934803

Option 4 ID : 41652934805

Status : Marked For Review



Chosen Option : 4

Q.3 The major product formed in the following reaction is :



Options

- 1.
- 2.



Question ID : 4165298820

Option 1 ID : 41652934738

Option 2 ID : 41652934740

Option 3 ID : 41652934739

Option 4 ID : 41652934741

Status : Not Answered

Chosen Option : --

Q.4 Which of the following conditions in drinking water causes methemoglobinemia ?

- Options
1. > 50 ppm of lead
  2. > 50 ppm of chloride
  3. > 50 ppm of nitrate
  4. > 100 ppm of sulphate

Question ID : 4165298834

Option 1 ID : 41652934795

Option 2 ID : 41652934797


Option 3 ID : 41652934796

Option 4 ID : 41652934794



Status : Marked For Review



Chosen Option : 2

Q.5 The major product of the following reaction is :



Options





Question ID : 4165298816

Option 1 ID : 41652934722

Option 2 ID : 41652934725


Option 3 ID : 41652934723

Option 4 ID : 41652934724





Status : Answered

Chosen Option : 3

Q.6 The major product obtained in the following reaction is :



Options



Question ID : 4165298824

Option 1 ID : 41652934756

Option 2 ID : 41652934757

Option 3 ID : 41652934754

Option 4 ID : 41652934755


Status : Not Answered

Chosen Option : --

Q.7



The major product of the following reaction is :



Options

- 1.
- 2.
- 3.
- 4.

Question ID : 4165298818

Option 1 ID : 41652934732

Option 2 ID : 41652934730

Option 3 ID : 41652934731

Option 4 ID : 41652934733

Status : Answered

Chosen Option : 3

Q.8 The correct match between Item I and Item II is :

Item I	Item II
(A) Benzaldehyde	(P) Mobile phase
(B) Alumina	(Q) Adsorbent
(C) Acetonitrile	(R) Adsorbate

- Options
1. (A) → (Q) ; (B) → (P) ; (C) → (R)
  2. (A) → (R) ; (B) → (Q) ; (C) → (P)
  3. (A) → (Q) ; (B) → (R) ; (C) → (P)
  4. (A) → (P) ; (B) → (R) ; (C) → (Q)

Question ID : 4165298825

Option 1 ID : 41652934759

Option 2 ID : 41652934761

Option 3 ID : 41652934760

Option 4 ID : 41652934758

Status : Answered

Chosen Option : 4

Q.9 The metal that forms nitride by reacting directly with  $N_2$  of air, is :

- Options 1. K

2. Li
3. Rb
4. Cs

Question ID : 4165298829  
Option 1 ID : 41652934774  
Option 2 ID : 41652934777  
Option 3 ID : 41652934775  
Option 4 ID : 41652934776  
Status : Answered  
Chosen Option : 1

Q.10 For coagulation of arsenious sulphide sol, which one of the following salt solution will be most effective ?

- Options
1.  $\text{BaCl}_2$
  2.  $\text{AlCl}_3$
  3.  $\text{NaCl}$
  4.  $\text{Na}_3\text{PO}_4$

Question ID : 4165298845  
Option 1 ID : 41652934839  
Option 2 ID : 41652934838  
Option 3 ID : 41652934840  
Option 4 ID : 41652934841  
Status : Not Answered  
Chosen Option : --

Q.11 The complex that has highest crystal field splitting energy ( $\Delta$ ), is :

- Options
1.  $[\text{Co}(\text{NH}_3)_5(\text{H}_2\text{O})]\text{Cl}_3$
  2.  $\text{K}_2[\text{CoCl}_4]$
  3.  $[\text{Co}(\text{NH}_3)_5\text{Cl}]\text{Cl}_2$
  4.  $\text{K}_3[\text{Co}(\text{CN})_6]$

Question ID : 4165298832  
Option 1 ID : 41652934789  
Option 2 ID : 41652934786  
Option 3 ID : 41652934788  
Option 4 ID : 41652934787  
Status : Answered  
Chosen Option : 4



Q.12 The pH of rain water, is approximately :

- Options
1. 5.6
  2. 7.5
  3. 7.0
  4. 6.5

Question ID : 4165298835  
Option 1 ID : 41652934801

Option 2 ID : 41652934799  
 Option 3 ID : 41652934798  
 Option 4 ID : 41652934800  
 Status : Answered  
 Chosen Option : 1

Q.13 Consider the following reversible chemical reactions :




The relation between  $K_1$  and  $K_2$  is :

- Options
1.  $K_1 K_2 = \frac{1}{3}$
  2.  $K_2 = K_1^3$
  3.  $K_2 = K_1^{-3}$
  4.  $K_1 K_2 = 3$

Question ID : 4165298842  
 Option 1 ID : 41652934829  
 Option 2 ID : 41652934827  
 Option 3 ID : 41652934826  
 Option 4 ID : 41652934828  
 Status : Answered  
 Chosen Option : 4

Q.14 The correct sequence of amino acids present in the tripeptide given below is :



- Options
1. Val - Ser - Thr
  2. Thr - Ser - Val
  3. Leu - Ser - Thr
  4. Thr - Ser - Leu

Question ID : 4165298822  
 Option 1 ID : 41652934748  
 Option 2 ID : 41652934749  
 Option 3 ID : 41652934746  
 Option 4 ID : 41652934747  
 Status : Not Answered  
 Chosen Option : --

Q.15

For the reaction,  $2A + B \rightarrow$  products, when the concentrations of A and B both were doubled, the rate of the reaction increased from  $0.3 \text{ mol L}^{-1}\text{s}^{-1}$  to  $2.4 \text{ mol L}^{-1}\text{s}^{-1}$ . When the concentration of A alone is doubled, the rate increased from  $0.3 \text{ mol L}^{-1}\text{s}^{-1}$  to  $0.6 \text{ mol L}^{-1}\text{s}^{-1}$ .

Which one of the following statements is correct ?

- Options
1. Total order of the reaction is 4
  2. Order of the reaction with respect to B is 2
  3. Order of the reaction with respect to B is 1
  4. Order of the reaction with respect to A is 2

Question ID : 4165298844

Option 1 ID : 41652934835

Option 2 ID : 41652934834

Option 3 ID : 41652934837





Option 4 ID : 41652934836

Status : Not Answered

Chosen Option : --

Q.16 The products formed in the reaction of cumene with  $\text{O}_2$  followed by treatment with dil. HCl are :

Options



Question ID : 4165298819

Option 1 ID : 41652934736

Option 2 ID : 41652934737

Option 3 ID : 41652934735

Option 4 ID : 41652934734

Status : Not Answered

Chosen Option : --





Q.17

The tests performed on compound X and their inferences are :

Test	Inference
(a) 2, 4 - DNP test	Coloured precipitate
(b) Iodoform test	Yellow precipitate
(c) Azo-dye test	No dye formation

Compound 'X' is :

Options



Question ID : 4165298823

Option 1 ID : 41652934752

Option 2 ID : 41652934751


Option 3 ID : 41652934750

Option 4 ID : 41652934753

Status : Not Answered

Chosen Option : --

Q.18 If the standard electrode potential for a cell is 2 V at 300 K, the equilibrium constant (K) for the reaction



at 300 K is approximately

( $R = 8 \text{ JK}^{-1}\text{mol}^{-1}$ ,  $F = 96000 \text{ C mol}^{-1}$ )

Options 1.  $e^{-80}$

2.  $e^{-160}$

3.  $e^{320}$

4.  $e^{160}$

Question ID : 4165298843

Option 1 ID : 41652934830

Option 2 ID : 41652934832

Option 3 ID : 41652934831

Option 4 ID : 41652934833

Status : **Marked For Review**  
Chosen Option : 4

Q.19 The temporary hardness of water is due to :

- Options
1.  $\text{Na}_2\text{SO}_4$
  2.  $\text{NaCl}$
  3.  $\text{Ca}(\text{HCO}_3)_2$
  4.  $\text{CaCl}_2$

Question ID : 4165298828  
Option 1 ID : 41652934773  
Option 2 ID : 41652934772  
Option 3 ID : 41652934771  
Option 4 ID : 41652934770  
Status : **Answered**  
Chosen Option : 4

Q.20 In which of the following processes, the bond order has increased and paramagnetic character has changed to diamagnetic ?

- Options
1.  $\text{NO} \rightarrow \text{NO}^+$
  2.  $\text{N}_2 \rightarrow \text{N}_2^+$
  3.  $\text{O}_2 \rightarrow \text{O}_2^+$
  4.  $\text{O}_2 \rightarrow \text{O}_2^{2-}$

Question ID : 4165298839  
Option 1 ID : 41652934814  
Option 2 ID : 41652934817  
Option 3 ID : 41652934815  
Option 4 ID : 41652934816  
Status : **Answered**  
Chosen Option : 1

Q.21 Which of the following combination of statements is true regarding the interpretation of the atomic orbitals ?

- (a) An electron in an orbital of high angular momentum stays away from the nucleus than an electron in the orbital of lower angular momentum.
- (b) For a given value of the principal quantum number, the size of the orbit is inversely proportional to the azimuthal quantum number.
- (c) According to wave mechanics, the ground state angular momentum is equal to  $\frac{h}{2\pi}$ .
- (d) The plot of  $\psi$  Vs  $r$  for various azimuthal quantum numbers, shows peak shifting towards higher  $r$  value.





- Options
1. (a), (d)

2. (a), (b)
3. (a), (c)
4. (b), (c)

Question ID : 4165298838  
 Option 1 ID : 41652934813  
 Option 2 ID : 41652934810  
 Option 3 ID : 41652934811  
 Option 4 ID : 41652934812  
 Status : Answered  
 Chosen Option : 4

Q.22 Which of the following compounds is not aromatic ?

Options

1. 
2. 
3. 
4. 

Question ID : 4165298817  
 Option 1 ID : 41652934728  
 Option 2 ID : 41652934727  
 Option 3 ID : 41652934726  
 Option 4 ID : 41652934729  
 Status : Answered  
 Chosen Option : 3


Q.23 Good reducing nature of  $\text{H}_3\text{PO}_2$  is attributed to the presence of :

- Options
1. Two P – OH bonds
  2. One P – H bond
  3. Two P – H bonds
  4. One P – OH bond

Question ID : 4165298830  
 Option 1 ID : 41652934781  
 Option 2 ID : 41652934778  
 Option 3 ID : 41652934779  
 Option 4 ID : 41652934780  
 Status : Answered  
 Chosen Option : 3

Q.24

The correct statement regarding the given Ellingham diagram is :



Options

1. At 1400°C, Al can be used for the extraction of Zn from ZnO
2. At 500°C, coke can be used for the extraction of Zn from ZnO
3. Coke cannot be used for the extraction of Cu from  $\text{Cu}_2\text{O}$ .
4. At 800°C, Cu can be used for the extraction of Zn from ZnO.

Question ID : 4165298827

Option 1 ID : 41652934767

Option 2 ID : 41652934768

Option 3 ID : 41652934766

Option 4 ID : 41652934769

Status : Not Answered

Chosen Option : --

Q.25 The transition element that has lowest enthalpy of atomisation, is :

- Options
1. Fe
  2. Cu
  3. V
  4. Zn

Question ID : 4165298831

Option 1 ID : 41652934785

Option 2 ID : 41652934783

Option 3 ID : 41652934782

Option 4 ID : 41652934784





Status : Answered

Chosen Option : 3

Q.26



The increasing basicity order of the following compounds is :



- Options
1. (D)<(C)<(B)<(A)
  2. (D)<(C)<(A)<(B)
  3. (A)<(B)<(C)<(D)
  4. (A)<(B)<(D)<(C)

Question ID : 4165298821  
 Option 1 ID : 41652934743  
 Option 2 ID : 41652934744  
 Option 3 ID : 41652934745  
 Option 4 ID : 41652934742  
 Status : Answered  
 Chosen Option : 1

Q.27 When the first electron gain enthalpy ( $\Delta_{\text{eg}}H$ ) of oxygen is  $-141 \text{ kJ/mol}$ , its second electron gain enthalpy is :

- Options
1. a more negative value than the first
  2. almost the same as that of the first
  3. negative, but less negative than the first
  4. a positive value

Question ID : 4165298826  
 Option 1 ID : 41652934763  
 Option 2 ID : 41652934765  
 Option 3 ID : 41652934764  
 Option 4 ID : 41652934762  
 Status : Answered  
 Chosen Option : 3

Q.28 At  $100^\circ\text{C}$ , copper (Cu) has FCC unit cell structure with cell edge length of  $x \text{ \AA}$ . What is the approximate density of Cu (in  $\text{g cm}^{-3}$ ) at this temperature ?  
 [Atomic Mass of Cu = 63.55 u]

- Options
1.  $\frac{205}{x^3}$
  2.  $\frac{105}{x^3}$

3.  $\frac{211}{x^3}$

4.  $\frac{422}{x^3}$

Question ID : 4165298837  
 Option 1 ID : 41652934809  
 Option 2 ID : 41652934806  
 Option 3 ID : 41652934807  
 Option 4 ID : 41652934808  
 Status : Not Answered  
 Chosen Option : --

Q.29 A solution containing 62 g ethylene glycol in 250 g water is cooled to  $-10^{\circ}\text{C}$ . If  $K_f$  for water is  $1.86 \text{ K kg mol}^{-1}$ , the amount of water (in g) separated as ice is :

- Options
1. 48
  2. 32
  3. 64
  4. 16

Question ID : 4165298841  
 Option 1 ID : 41652934824  
 Option 2 ID : 41652934823  
 Option 3 ID : 41652934825  
 Option 4 ID : 41652934822  
 Status : Marked For Review  
 Chosen Option : 4

Q.30 Homoleptic octahedral complexes of a metal ion ' $M^{3+}$ ' with three monodentate ligands  $L_1$ ,  $L_2$  and  $L_3$  absorb wavelengths in the region of green, blue and red respectively. The increasing order of the ligand strength is :

- Options
1.  $L_3 < L_1 < L_2$
  2.  $L_3 < L_2 < L_1$
  3.  $L_1 < L_2 < L_3$
  4.  $L_2 < L_1 < L_3$

Question ID : 4165298833  
 Option 1 ID : 41652934793  
 Option 2 ID : 41652934791  
 Option 3 ID : 41652934790  
 Option 4 ID : 41652934792  
 Status : Answered  
 Chosen Option : 4

Section : Mathematics

Q.1

The sum of the following series

$$1+6+\frac{9(1^2+2^2+3^2)}{7}+\frac{12(1^2+2^2+3^2+4^2)}{9} \\ +\frac{15(1^2+2^2+\dots+5^2)}{11}+\dots \text{ up to 15 terms,}$$

is :

- Options
1. 7520
  2. 7510
  3. 7830
  4. 7820

Question ID : 4165298854

Option 1 ID : 41652934877

Option 2 ID : 41652934874

Option 3 ID : 41652934876

Option 4 ID : 41652934875

Status : Not Answered

Chosen Option : --

Q.2 For each  $x \in \mathbf{R}$ , let  $[x]$  be the greatest integer less than or equal to  $x$ . Then

$$\lim_{x \rightarrow 0^-} \frac{x([x]+|x|) \sin [x]}{|x|} \text{ is equal to :}$$

- Options
1.  $-\sin 1$
  2. 1
  3.  $\sin 1$
  4. 0

Question ID : 4165298855

Option 1 ID : 41652934880

Option 2 ID : 41652934881

Option 3 ID : 41652934878

Option 4 ID : 41652934879

Status : Answered

Chosen Option : 4

Q.3 Let  $f: [0, 1] \rightarrow \mathbf{R}$  be such that  $f(xy) = f(x) \cdot f(y)$ , for all  $x, y \in [0, 1]$ , and  $f(0) \neq 0$ . If  $y = y(x)$  satisfies the differential equation,

$$\frac{dy}{dx} = f(x) \text{ with } y(0) = 1, \text{ then } y\left(\frac{1}{4}\right) + y\left(\frac{3}{4}\right)$$

is equal to :

- Options
1. 3
  2. 4
  3. 2
  4. 5

Question ID : 4165298862

Option 1 ID : 41652934907

Option 2 ID : 41652934908  
 Option 3 ID : 41652934906  
 Option 4 ID : 41652934909  
 Status : Marked For Review  
 Chosen Option : 2

Q.4 If  $x = \sin^{-1}(\sin 10)$  and  $y = \cos^{-1}(\cos 10)$ ,  
 then  $y - x$  is equal to :

- Options
1. 0
  2. 10
  3.  $7\pi$
  4.  $\pi$

Question ID : 4165298874  
 Option 1 ID : 41652934954  
 Option 2 ID : 41652934956  
 Option 3 ID : 41652934957  
 Option 4 ID : 41652934955  
 Status : Answered  
 Chosen Option : 1

Q.5 If  $0 \leq x < \frac{\pi}{2}$ , then the number of values of  $x$   
 for which  $\sin x - \sin 2x + \sin 3x = 0$ , is :

- Options
1. 3
  2. 1
  3. 4
  4. 2

Question ID : 4165298873  
 Option 1 ID : 41652934950  
 Option 2 ID : 41652934953  
 Option 3 ID : 41652934951  
 Option 4 ID : 41652934952  
 Status : Answered  
 Chosen Option : 4

Q.6 Let  $z_0$  be a root of the quadratic equation,  
 $x^2 + x + 1 = 0$ . If  $z = 3 + 6iz_0^{81} - 3iz_0^{93}$ , then  
 arg  $z$  is equal to :

- Options
1.  $\frac{\pi}{4}$
  2.  $\frac{\pi}{6}$
  3.  $\frac{\pi}{3}$
  4. 0

Question ID : 4165298847  
 Option 1 ID : 41652934846  
 Option 2 ID : 41652934847  
 Option 3 ID : 41652934848  
 Option 4 ID : 41652934849

Status : **Not Answered**  
Chosen Option : --

Q.7 The area of the region

$A = \{(x, y) : 0 \leq y \leq x|x| + 1 \text{ and } -1 \leq x \leq 1\}$  in sq. units, is :

- Options
1.  $\frac{2}{3}$
  2. 2
  3.  $\frac{4}{3}$
  4.  $\frac{1}{3}$

Question ID : 4165298861  
Option 1 ID : 41652934902  
Option 2 ID : 41652934905  
Option 3 ID : 41652934904  
Option 4 ID : 41652934903  
Status : **Answered**  
Chosen Option : 2

Q.8 If the system of linear equations

$$x - 4y + 7z = g$$

$$3y - 5z = h$$

$$-2x + 5y - 9z = k$$

is consistent, then :

- Options
1.  $g + 2h + k = 0$
  2.  $g + h + 2k = 0$
  3.  $2g + h + k = 0$
  4.  $g + h + k = 0$

Question ID : 4165298850  
Option 1 ID : 41652934859  
Option 2 ID : 41652934860  
Option 3 ID : 41652934858  
Option 4 ID : 41652934861  
Status : **Answered**  
Chosen Option : 3

Q.9 The coefficient of  $t^4$  in the expansion of

$$\left( \frac{1-t^6}{1-t} \right)^3 \text{ is :}$$

- Options
1. 14
  2. 15
  3. 10
  4. 12

Question ID : 4165298852  
Option 1 ID : 41652934869

Option 2 ID : 41652934868  
 Option 3 ID : 41652934867  
 Option 4 ID : 41652934866  
 Status : Marked For Review  
 Chosen Option : 4

Q.10 If both the roots of the quadratic equation  $x^2 - mx + 4 = 0$  are real and distinct and they lie in the interval  $[1, 5]$ , then  $m$  lies in the interval :

- Options
1.  $(-5, -4)$
  2.  $(4, 5)$
  3.  $(5, 6)$
  4.  $(3, 4)$

Question ID : 4165298857  
 Option 1 ID : 41652934886  
 Option 2 ID : 41652934888  
 Option 3 ID : 41652934889  
 Option 4 ID : 41652934887  
 Status : Marked For Review  
 Chosen Option : 4

Q.11 Let  $S$  be the set of all triangles in the  $xy$ -plane, each having one vertex at the origin and the other two vertices lie on coordinate axes with integral coordinates. If each triangle in  $S$  has area 50 sq. units, then the number of elements in the set  $S$  is :

- Options
1. 9
  2. 18
  3. 36
  4. 32

Question ID : 4165298863  
 Option 1 ID : 41652934910  
 Option 2 ID : 41652934911  
 Option 3 ID : 41652934912  
 Option 4 ID : 41652934913  
 Status : Answered  
 Chosen Option : 3

Q.12 Let  $a, b$  and  $c$  be the 7<sup>th</sup>, 11<sup>th</sup> and 13<sup>th</sup> terms respectively of a non-constant A.P. If these are also the three consecutive terms of a G.P., then  $\frac{a}{c}$  is equal to :

- Options
1. 2
  2.  $\frac{1}{2}$
  3.  $\frac{7}{13}$

4. 4

Question ID : 4165298853  
 Option 1 ID : 41652934871  
 Option 2 ID : 41652934873  
 Option 3 ID : 41652934872  
 Option 4 ID : 41652934870  
 Status : Marked For Review  
 Chosen Option : 1

Q.13 The logical statement  
 $[\sim(\sim p \vee q) \vee (p \wedge r)] \wedge (\sim q \wedge r)$   
 is equivalent to :

- Options
1.  $(\sim p \wedge \sim q) \wedge r$
  2.  $\sim p \vee r$
  3.  $(p \wedge r) \wedge \sim q$
  4.  $(p \wedge \sim q) \vee r$

Question ID : 4165298875  
 Option 1 ID : 41652934961  
 Option 2 ID : 41652934958  
 Option 3 ID : 41652934959  
 Option 4 ID : 41652934960  
 Status : Not Answered  
 Chosen Option : --

Q.14 The equation of the plane containing the  
 straight line  $\frac{x}{2} = \frac{y}{3} = \frac{z}{4}$  and perpendicular  
 to the plane containing the straight lines

$$\frac{x}{3} = \frac{y}{4} = \frac{z}{2} \text{ and } \frac{x}{4} = \frac{y}{2} = \frac{z}{3} \text{ is :}$$

- Options
1.  $x - 2y + z = 0$
  2.  $3x + 2y - 3z = 0$
  3.  $x + 2y - 2z = 0$
  4.  $5x + 2y - 4z = 0$

Question ID : 4165298869  
 Option 1 ID : 41652934936  
 Option 2 ID : 41652934935  
 Option 3 ID : 41652934934  
 Option 4 ID : 41652934937  
 Status : Answered  
 Chosen Option : 1

Q.15 A data consists of n observations :

$$x_1, x_2, \dots, x_n. \text{ If } \sum_{i=1}^n (x_i + 1)^2 = 9n \text{ and}$$

$$\sum_{i=1}^n (x_i - 1)^2 = 5n, \text{ then the standard}$$

deviation of this data is :

Options 1. 2

2.  $\sqrt{5}$

3. 5

4.  $\sqrt{7}$

Question ID : 4165298871

Option 1 ID : 41652934945

Option 2 ID : 41652934943

Option 3 ID : 41652934944

Option 4 ID : 41652934942

Status : Not Answered

Chosen Option : --

Q.16 If

$$A = \begin{bmatrix} e^t & e^{-t} \cos t & e^{-t} \sin t \\ e^t & -e^{-t} \cos t - e^{-t} \sin t & -e^{-t} \sin t + e^{-t} \cos t \\ e^t & 2e^{-t} \sin t & -2e^{-t} \cos t \end{bmatrix},$$

then A is :

Options 1. invertible for all  $t \in \mathbb{R}$ .2. invertible only if  $t = \pi$ .3. not invertible for any  $t \in \mathbb{R}$ .4. invertible only if  $t = \frac{\pi}{2}$ .

Question ID : 4165298849

Option 1 ID : 41652934856

Option 2 ID : 41652934854

Option 3 ID : 41652934857

Option 4 ID : 41652934855

Status : Answered

Chosen Option : 1

Q.17

If  $f(x) = \int \frac{5x^8 + 7x^6}{(x^2 + 1 + 2x^7)^2} dx, (x \geq 0)$ , and

 $f(0) = 0$ , then the value of  $f(1)$  is :

Options 1.  $-\frac{1}{2}$

2.  $-\frac{1}{4}$

3.  $\frac{1}{2}$

4.  $\frac{1}{4}$

Question ID : 4165298859

Option 1 ID : 41652934894

Option 2 ID : 41652934897



Option 3 ID : 41652934896

Option 4 ID : 41652934895

Status : Answered

Chosen Option : 4

Q.18 Let  $f$  be a differentiable function from  $\mathbf{R}$  to  $\mathbf{R}$  such that  $|f(x) - f(y)| \leq 2|x - y|^{3/2}$ , for all  $x, y \in \mathbf{R}$ . If  $f(0) = 1$  then  $\int_0^1 f^2(x) dx$  is equal to :

Options 1. 1

2. 2

3.  $\frac{1}{2}$ 

4. 0

Question ID : 4165298858

Option 1 ID : 41652934891

Option 2 ID : 41652934893

Option 3 ID : 41652934892

Option 4 ID : 41652934890

Status : Answered

Chosen Option : 2

Q.19 If  $x = 3 \tan t$  and  $y = 3 \sec t$ , then the value of  $\frac{d^2 y}{dx^2}$  at  $t = \frac{\pi}{4}$ , is :

Options 1.  $\frac{1}{3\sqrt{2}}$ 2.  $\frac{1}{6\sqrt{2}}$ 3.  $\frac{3}{2\sqrt{2}}$ 4.  $\frac{1}{6}$ 

Question ID : 4165298856

Option 1 ID : 41652934884

Option 2 ID : 41652934883

Option 3 ID : 41652934885

Option 4 ID : 41652934882

Status : Answered

Chosen Option : 1

Q.20 The number of natural numbers less than 7,000 which can be formed by using the digits 0, 1, 3, 7, 9 (repetition of digits allowed) is equal to :

Options 1. 374

2. 372

3. 375

4. 250

Question ID : 4165298851

Option 1 ID : 41652934864

Option 2 ID : 41652934862

Option 3 ID : 41652934863

Option 4 ID : 41652934865

Status : Answered

Chosen Option : 3

Q.21 If the circles  $x^2 + y^2 - 16x - 20y + 164 = r^2$  and  $(x - 4)^2 + (y - 7)^2 = 36$  intersect at two distinct points, then :

Options 1.  $r > 11$ 2.  $0 < r < 1$ 3.  $r = 11$ 4.  $1 < r < 11$ 

Question ID : 4165298865

Option 1 ID : 41652934921

Option 2 ID : 41652934919

Option 3 ID : 41652934920

Option 4 ID : 41652934918

Status : Marked For Review

Chosen Option : 3

Q.22 A hyperbola has its centre at the origin, passes through the point (4, 2) and has transverse axis of length 4 along the x-axis. Then the eccentricity of the hyperbola is :

Options 1.  $\frac{3}{2}$ 2.  $\sqrt{3}$ 

3. 2

4.  $\frac{2}{\sqrt{3}}$ 

Question ID : 4165298867

Option 1 ID : 41652934929

Option 2 ID : 41652934926

Option 3 ID : 41652934928

Option 4 ID : 41652934927

Status : Answered

Chosen Option : 1

Q.23 Let A(4, -4) and B(9, 6) be points on the parabola,  $y^2 = 4x$ . Let C be chosen on the arc AOB of the parabola, where O is the origin, such that the area of  $\Delta ACB$  is maximum. Then, the area (in sq.units) of  $\Delta ACB$ , is :

Options

1.  $31\frac{1}{4}$
2.  $30\frac{1}{2}$
3. 32
4.  $31\frac{3}{4}$

Question ID : 4165298866  
 Option 1 ID : 41652934922  
 Option 2 ID : 41652934923  
 Option 3 ID : 41652934924  
 Option 4 ID : 41652934925  
 Status : **Marked For Review**  
 Chosen Option : 3

**Q.24** Let the equations of two sides of a triangle be  $3x - 2y + 6 = 0$  and  $4x + 5y - 20 = 0$ . If the orthocentre of this triangle is at  $(1, 1)$ , then the equation of its third side is :

- Options
1.  $122y - 26x - 1675 = 0$
  2.  $122y + 26x + 1675 = 0$
  3.  $26x + 61y + 1675 = 0$
  4.  $26x - 122y - 1675 = 0$

Question ID : 4165298864  
 Option 1 ID : 41652934914  
 Option 2 ID : 41652934917  
 Option 3 ID : 41652934916  
 Option 4 ID : 41652934915  
 Status : **Not Answered**  
 Chosen Option : --

**Q.25** An urn contains 5 red and 2 green balls. A ball is drawn at random from the urn. If the drawn ball is green, then a red ball is added to the urn and if the drawn ball is red, then a green ball is added to the urn ; the original ball is not returned to the urn. Now, a second ball is drawn at random from it. The probability that the second ball is red, is :

- Options
1.  $\frac{21}{49}$
  2.  $\frac{27}{49}$
  3.  $\frac{26}{49}$
  4.  $\frac{32}{49}$

Question ID : 4165298872

Option 1 ID : 41652934947  
 Option 2 ID : 41652934946  
 Option 3 ID : 41652934948  
 Option 4 ID : 41652934949  
 Status : Answered  
 Chosen Option : 1

Q.26 If the lines  $x = ay + b$ ,  $z = cy + d$  and  $x = a'z + b'$ ,  $y = c'z + d'$  are perpendicular, then :

- Options
1.  $ab' + bc' + 1 = 0$
  2.  $cc' + a + a' = 0$
  3.  $bb' + cc' + 1 = 0$
  4.  $aa' + c + c' = 0$

Question ID : 4165298868  
 Option 1 ID : 41652934930  
 Option 2 ID : 41652934933  
 Option 3 ID : 41652934931  
 Option 4 ID : 41652934932  
 Status : Not Answered  
 Chosen Option : --

Q.27 Let  $\vec{a} = \hat{i} + \hat{j} + \sqrt{2}\hat{k}$ ,  $\vec{b} = b_1\hat{i} + b_2\hat{j} + \sqrt{2}\hat{k}$  and  $\vec{c} = 5\hat{i} + \hat{j} + \sqrt{2}\hat{k}$  be three vectors such that the projection vector of  $\vec{b}$  on  $\vec{a}$  is  $\vec{a}$ . If  $\vec{a} + \vec{b}$  is perpendicular to  $\vec{c}$ , then  $|\vec{b}|$  is equal to :

- Options
1.  $\sqrt{32}$
  2. 6
  3.  $\sqrt{22}$
  4. 4

Question ID : 4165298870  
 Option 1 ID : 41652934941  
 Option 2 ID : 41652934938  
 Option 3 ID : 41652934940  
 Option 4 ID : 41652934939  
 Status : Marked For Review  
 Chosen Option : 4

Q.28 The number of all possible positive integral values of  $\alpha$  for which the roots of the quadratic equation,  $6x^2 - 11x + \alpha = 0$  are rational numbers is :

- Options
1. 3
  2. 2
  3. 4
  4. 5

Question ID : 4165298848

Option 1 ID : 41652934851

Option 2 ID : 41652934850

Option 3 ID : 41652934852

Option 4 ID : 41652934853

Status : Answered

Chosen Option : 3

Q.29 Let  $A = \{x \in \mathbb{R} : x \text{ is not a positive integer}\}$ .

Define a function  $f: A \rightarrow \mathbb{R}$  as  $f(x) = \frac{2x}{x-1}$ ,

then  $f$  is :

- Options
1. not injective
  2. neither injective nor surjective
  3. surjective but not injective
  4. injective but not surjective

Question ID : 4165298846

Option 1 ID : 41652934842

Option 2 ID : 41652934845

Option 3 ID : 41652934844

Option 4 ID : 41652934843

Status : Answered

Chosen Option : 4

Q.30

If  $\int_0^{\pi/3} \frac{\tan \theta}{\sqrt{2k \sec \theta}} d\theta = 1 - \frac{1}{\sqrt{2}}$ , ( $k > 0$ ), then the

value of  $k$  is :

- Options
1. 4
  2.  $\frac{1}{2}$
  3. 1
  4. 2

Question ID : 4165298860

Option 1 ID : 41652934900

Option 2 ID : 41652934901

Option 3 ID : 41652934899

Option 4 ID : 41652934898

Status : Answered

Chosen Option : 2