

ALCOHOL, PHENOL, ETHER





(ORGANIC CHAPTER-2)

ALCOHOL

IUPAC name:- Alkanol

Common name:- Alkyl alcohol




IUPAC names of some alcohols that were asked in board examination:-

STRUCTURAL FORMULA	IUPAC name
1. $\begin{array}{c} \text{CH}_3 - \text{C} = \text{C} - \text{CH}_2 - \text{OH} \\ \quad \\ \text{CH}_3 \quad \text{Br} \end{array}$	2-bromo-3-methylbut-2-en-1-ol
2. 	3-phenylprop-2-en-1-ol
3. $\begin{array}{c} \text{OH} - \text{CH}_2 - \text{CH} - \text{CH}_2 - \text{OH} \\ \\ \text{CH}_3 \end{array}$	2-methylpropane-1,3-diol
4. 	2,6-dimethylphenol
5. $\begin{array}{c} \text{CH}_3 - \text{O} - \text{CH}_2 - \text{CH} - \text{CH}_3 \\ \\ \text{OH} \end{array}$	1-methoxypropan-2-ol
6. 	2-phenylethanol
7. 	2,5-dinitrophenol
8. $\begin{array}{c} \text{OH} - \text{CH}_2 - \text{CH} = \text{C} - \text{CH}_3 \\ \\ \text{CH}_3 \end{array}$	3-methylbut-2-en-1-ol

PREPARATION OF ALCOHOLS:-


① FROM ALKENES:-

(i) By acid catalysed hydration:-




Mechanism:-


Step-1:- Formation of carbocation




Step-2:- Nucleophilic attack of water on carbocation




Step-3:- Deprotonation to form an alcohol




(ii) By hydroboration oxidation:-




(Here, water is added by anti-markovnikov's rule)




In this process, primary alcohol is formed.



2 FROM CARBONYL COMPOUND:-





Eg:-





(ii) Reduction of aldehydes and ketenes:-

In Aldehyde,





In Ketone,






(iii) Reduction of carboxylic Acid and ester:-

In carboxylic acid,




In ester,











4

* NaBH_4 cannot reduce ester whereas LiAlH_4 can.




③ FROM GRIGNARD REAGENTS:-




REPARATION OF PHENOL:-

① From haloarenes / chlorobenzene:-




JM


② From benzene sulphonic acid:-



③ From diazonium salt:-



④ From Cumene:-



PHYSICAL PROPERTIES:-

(i) BOILING POINT:- More the number of C-atoms in the straight chain, more surface area, more vanderwall force so more boiling point. The boiling point decreases with branching.

(ii) SOLUBILITY:- Alcohols are soluble in water due to intermolecular hydrogen bonding. Lower alcohols are soluble in water. On increasing the number of carbon chain, due to +I effect of alkyl group, the polarity of OH decreases, so the tendency to form hydrogen bond decreases. So higher alcohols are less soluble or insoluble in water.


CHEMICAL PROPERTIES:-

Alcohols have two types of bond breaking reaction.

- (a) C-O bond breaking
- (b) O-H bond breaking

* C-O bond of alcohol will break when it undergoes nucleophilic substitution reaction. So the bond between C-O is broken when they react as electrophiles.



* The bond between O-H is broken when alcohols react as nucleophiles.



(a) Reactions involving cleavage of O-H bond:-

1. Acidity of alcohols and phenols:-


(i) Reaction with metals:-




(ii) Acidic nature of alcohol:-

- Electron withdrawing group (NO_2 , CN , Aldehyde, Ketone, carboxylic acid, halogen) increases the acidic character.
- Electron donating group (Alkyl, ether, alcohol) decreases the acidic character.

(iii) Acidity of phenols:-



In case of phenol, due to resonance charge separation is there. So kinetically, its energy is increased which leads to lesser stability.





In case of phenoxide ion, only negative charge is delocalised, so there is no charge separation. So, phenoxide ion is more stable.

(b) Reactions involving cleavage of carbon-oxygen bond in alcohols:- ⑦


① Dehydration of alcohol:-

Here, conc H_2SO_4 or conc H_3PO_4 act as dehydrating agent.




Mechanism:-

① Formation of protonated alcohol





② Formation of carbocation





③ Deprotonation



Secondary and Tertiary alcohols are dehydrated under milder conditions. For example:-




The relative ease of dehydration of alcohols follows the following order:- Tertiary > Secondary > Primary.




(c) Reaction of phenols:-

1. Electrophilic substitution reaction:-




As, negative charge is delocalised at ortho and para position. So it is a 'strong activating group'.


(i) Nitration:-




The ortho and para isomers can be separated by steam distillation. o-Nitrophenol is steam volatile due to intramolecular hydrogen bonding while p-Nitrophenol is less volatile due to intermolecular hydrogen bonding.




(ii) Halogenation:-







② Oxidation of phenol:-




③ Reaction with zinc dust:-



④ Reimer-Tiemann reaction:- (Phenol \rightarrow salicylaldehyde)



⑤ Kolbe's reaction:- (Phenol \rightarrow salicylic acid)




The commercial alcohol is made unfit for drinking by mixing in it some copper sulphate (to give colour) & pyridine (a foul smelling liquid). It is known as denaturation of alcohol.

PREPARATION OF ETHERS :-


① Dehydration of alcohol :-

At 413 K, two OH groups elimination of water takes place.




MECHANISM:-



Step-①:- Protonation.




Step-②:- Formation of carbocation.



Step-③:- Deprotonation




② WILLIAMSON SYNTHESIS:-



Alkyl
Halide

Eg:-




Better results are obtained if alkyl halide is primary. In case of 2° or 3° alkyl halide, elimination predominates over substitution. In case of 3° alkyl halide, alkene is formed instead of ether.

PHYSICAL PROPERTIES:-


Boiling Point :- Carboxylic acid > Alcohol > Aldehyde > Ketone > Ether > Alkane

CHEMICAL PROPERTIES:-


Case-I:- If alkyl group is 1° or 2°




Case-II:- If alkyl group is 3°




Case-III:- If one side of alkyl group is phenyl




Case-IV:- If one side is benzyl or allyl




② ELECTROPHILIC SUBSTITUTION REACTION OF ANISOLE:-



② FRIEDEL-CRAFT'S ALKYLATION:-



③ FRIEDEL-CRAFT'S ACYLATION:-



④ NITRATION:-

